

Outer Length Scales in Nocturnal Stable Boundary Layers

Basu, Sukanta

DOI

[10.1364/PCAOP.2022.PTu3D.3](https://doi.org/10.1364/PCAOP.2022.PTu3D.3)

Publication date

2022

Document Version

Final published version

Published in

Propagation Through and Characterization of Atmospheric and Oceanic Phenomena, pcAOP 2022

Citation (APA)

Basu, S. (2022). Outer Length Scales in Nocturnal Stable Boundary Layers. In *Propagation Through and Characterization of Atmospheric and Oceanic Phenomena, pcAOP 2022* (pp. 1-2). Article PTu3D.3 (Optics InfoBase Conference Papers). Optica Publishing Group (formerly OSA).
<https://doi.org/10.1364/PCAOP.2022.PTu3D.3>

Important note

To cite this publication, please use the final published version (if applicable).
Please check the document version above.

Copyright

Other than for strictly personal use, it is not permitted to download, forward or distribute the text or part of it, without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license such as Creative Commons.

Takedown policy

Please contact us and provide details if you believe this document breaches copyrights.
We will remove access to the work immediately and investigate your claim.

Green Open Access added to TU Delft Institutional Repository

'You share, we take care!' - Taverne project

<https://www.openaccess.nl/en/you-share-we-take-care>

Otherwise as indicated in the copyright section: the publisher is the copyright holder of this work and the author uses the Dutch legislation to make this work public.

Outer Length Scales in Nocturnal Stable Boundary Layers

Sukanta Basu

*Faculty of Civil Engineering and Geosciences, Delft University of Technology, Delft, the Netherlands
s.basu@tudelft.nl*

Abstract: Recently, Basu and Holstlag (2021) proposed a unified framework for describing outer length scales (OLS). By utilizing this framework, we document various characteristics of OLS in nocturnal boundary layers over the US Great Plains. © 2022 The Author(s)

OCIS codes: (000.0010) Atmospheric and Oceanic Optics; (010.1330) Atmospheric Turbulence

Based on the variance and flux budget equations, Basu and Holstlag [1] derived closed-form solutions for outer length scale (OLS; denoted as L_X) and turbulent Prandtl number (Pr_t) for steady-state, stably stratified conditions. Specifically, they deduced:

$$L_X = \left(\frac{\sqrt{Pr_{t0} Pr_t}}{c_\theta} \right) \left(\frac{\sigma_\theta}{\Gamma} \right), \quad (1)$$

where the standard deviation of potential temperature is σ_θ . The gradient of mean potential temperature is represented by Γ . The turbulent Prandtl number for non-buoyant flows is denoted by Pr_{t0} ; it is typically assumed to be equal to 0.85. The coefficient c_θ is approximately equal to 2. This newly proposed OLS (L_X) was shown to be related to several other well-known characteristic length scales of turbulence (e.g., Hunt length scale, Ellison length scale, Ozmidov length scale) for different asymptotic stability conditions (e.g., near-neutral, very stable). Furthermore, various analytical results of [1] were in close agreement with published observational and direct numerical simulation generated data (e.g., [2, 3]).

According to the hypothesis by Kolmogorov–Obukhov–Corrsin, within the inertial-convective range (r), the second-order structure function of potential temperature (S_2^T) is written as:

$$S_2^T(r) = C_T^2 r^{2/3}, \quad (2)$$

where C_T^2 is the so-called temperature structure parameter. Based on the results from [1], Basu and Holstlag [4] further derived:

$$C_T^2 = \left(\frac{c Pr_{t0}}{c_\theta^2} \right) \left(\frac{\sigma_\theta^2}{L_X^{2/3}} \right), \quad (3)$$

where the coefficient c is typically assumed to be around 3.2. By plugging in the typical values of the various coefficients, we can simplify Eq. (3) as follows:

$$C_T^2 = c_X \left(\frac{\sigma_\theta^2}{L_X^{2/3}} \right), \quad (4)$$

where, c_X is approximately equal to 0.68.

Under the assumptions of stationarity and homogeneity, the following relationship can be easily derived from the definition of S_2^T :

$$S_2^T(r) = 2\sigma_\theta^2 [1 - C(r)], \quad (5)$$

where $C(r)$ is the autocorrelation function of potential temperature. By combining Eqs. (2), (4), and (5), one can arrive at:

$$C(r) = 1 - \left(\frac{c_X}{2} \right) \left(\frac{r}{L_X} \right)^{2/3}. \quad (6)$$

Thus, for $r = L_X$, the autocorrelation is approximately equal to 0.66. This simple finding is rather powerful as it will allow one to estimate L_X solely from measured temperature time series. Furthermore, with the estimated value of L_X , Eq. (4) can be subsequently used to predict the associated C_T^2 value.

During this presentation, we will demonstrate the prowess of the proposed approach by using measurement data from the CASES-99 field campaign [6]. For our analyses, data from sonic anemometers located at seven levels (5, 10, 20, 30, 40, 50, and 55 m) on a 60-m tower are considered. A few examples are shown in Fig. 1.




Fig. 1. Structure function analysis of four representative (normalized) temperature (θ_n) time series from the CASES-99 field campaign. Each time series is ten minutes long (sampling frequency of 20 Hz). The normalized series are shown at the bottom of the corresponding structure function plots. The black circles denote measured S_2^T values. The magenta colored lines on the structure function plots denote the estimated L_X values. The estimated C_T^2 values and other meteorological variables are also reported on these plots. From the estimated C_T^2 values, one can predict S_2^T values by making use of Eq. (2). These predicted S_2^T values are depicted on the structure function plots as red lines. The vertical blue lines simply denote $r = z$, where z is the height of the sonic anemometers. The horizontal blue lines represent $S_2^T = 2$. When the autocorrelation drops to zero, the ratio (S_2^T / σ_θ^2) approaches to 2 according to Eq. (5).

It is important to note that, more than fifty years ago, Fried [5] proposed a similar (not the same) formulation for OLS estimation from the autocorrelation function. In contrast to our analytical approach (utilizing the variance and flux budget equations), Fried's approach was based on heuristic arguments.

References

1. S. Basu, and A. A. M. Holtslag (2021) Turbulent Prandtl number and characteristic length scales in stably stratified flows: steady-state analytical solutions. *Environ Fluid Mech* 21:1273–1302
2. S. Basu, P. He, and A. W. DeMarco (2021a) Parameterizing the energy dissipation rate in stably stratified flows. *Boundary-Layer Meteorol* 178: 167–184
3. S. Basu, A. W. DeMarco, and P. He (2021b) On the dissipation rate of temperature fluctuations in stably stratified flows. *Environ Fluid Mech* 21: 63–82
4. S. Basu, and A. A. M. Holtslag (2022) Revisiting and revising Tatarskii's formulation for the temperature structure parameter (C_T^2) in atmospheric flows. *Environ Fluid Mech* 10.1007/s10652-022-09880-3
5. D. L. Fried (1967) Optical heterodyne detection of an atmospherically distorted signal wave front. *Proceedings of the IEEE* 55: 57–67
6. G. S. Poulos, and Coauthors (2002) CASES-99: a comprehensive investigation of the stable nocturnal boundary layer. *Bull Amer Meteorol Soc* 83: 555–581
7. V. I. Tatarskii (1971) The effects of the turbulent atmosphere on wave propagation. Israel Program for Scientific Translations, 472 pp