

Existence of solutions to the generalized periodic fractional boundary value problem

Fečkan, Michal; Marynets, Kateryna; Wang, Jin Rong

DOI 10.1002/mma.9097

Publication date 2023 **Document Version**

Final published version

Published in Mathematical Methods in the Applied Sciences

Citation (APA) Fečkan, M., Marynets, K., & Wang, J. R. (2023). Existence of solutions to the generalized periodic fractional boundary value problem. *Mathematical Methods in the Applied Sciences*, *46*(11), 11971-11982. https://doi.org/10.1002/mma.9097

Important note

To cite this publication, please use the final published version (if applicable). Please check the document version above.

Copyright

Other than for strictly personal use, it is not permitted to download, forward or distribute the text or part of it, without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license such as Creative Commons.

Takedown policy

Please contact us and provide details if you believe this document breaches copyrights. We will remove access to the work immediately and investigate your claim.

SPECIAL ISSUE PAPER

WILEY

Existence of solutions to the generalized periodic fractional boundary value problem

Michal Fečkan^{1,2} | Kateryna Marynets³ | JinRong Wang⁴

¹Department of Mathematical Analysis and Numerical Mathematics, Comenius University in Bratislava, Bratislava, Slovakia

²Mathematical Institute of Slovak Academy of Sciences, Bratislava, Slovakia

³Delft Institute of Applied Mathematics, Faculty of Electrical Engineering, Mathematics and Computer Science, Delft University of Technology, Delft, The Netherlands

⁴Department of Mathematics, Guizhou University, Guiyang, China

Correspondence

Kateryna Marynets, Delft Institute of Applied Mathematics, Faculty of Electrical Engineering, Mathematics and Computer Science, Delft University of Technology, Mekelweg 4, 2628CD Delft, The Netherlands. Email: K.Marynets@tudelft.nl

Communicated by: W. Szymańska-Dbowska

Funding information

This work is partially supported by the National Natural Science Foundation of China (11661016), Training Object of High Level and Innovative Talents of Guizhou Province ((2016)4006), Major Research Project of Innovative Group in Guizhou Education Department ([2018]012), the Slovak Research and Development Agency under the contract No. APVV-18-0308, and the Slovak Grant Agency VEGA No. 1/0084/23 and No. 2/0127/20. We study a boundary value problem for a Caputo-type fractional differential equation subjected to periodic boundary conditions. For an auxiliary problem with the simplified right-hand side, we explicitly construct its unique solution. In addition, based on the theory of the topological index, we prove existence of at least one solution to the original problem.

KEYWORDS

existence of solutions, fractional boundary value problem, periodic boundary conditions, topological index

MSC CLASSIFICATION

34A08

1 | INTRODUCTION

In the theory of applied fractional calculus, boundary value problems for $periodic^{1-6}$ and $anti-periodic^{7-10}$ boundary conditions play an important role. They often occur in mathematical models of the real-world problems, for example, in

This is an open access article under the terms of the Creative Commons Attribution-NonCommercial-NoDerivs License, which permits use and distribution in any medium, provided the original work is properly cited, the use is non-commercial and no modifications or adaptations are made. © 2023 The Authors. Mathematical Methods in the Applied Sciences published by John Wiley & Sons Ltd.

epidemiology, ecology, physics, and material sciences (see Kilbas et al¹¹ and Podbulny¹²). This motivates the in-depth study of these types of problems aiming to prove existence and/or uniqueness of their solutions.

Among the already known results in study of the periodic FBVPs including the construction of an explicit solution are those obtained in previous studies.^{3–5} For instance, Fečkan and Marynets⁴ study the BVP

$${}_{0}^{C}D_{t}^{p}x(t) = f(t, x(t)), \ p \in (0, 1),$$
(1)

$$x(0) = x(T), \tag{2}$$

where ${}_{0}^{C}D_{t}^{p}$ is the generalized Caputo fractional derivative with lower limit at 0 (see Podbulny¹² and Zhou¹²), $t \in [0, T], x : [0, T] \to D$, $f : G \to \mathbb{R}^{n}$ are continuous functions, $G := [0, T] \times D$ and $D \subset \mathbb{R}^{n}$ is a closed and bounded domain. Under assumptions that function f in the system (1) is bounded by a constant vector $M = (M_{1}, M_{2}, \dots, M_{n})^{T} \in \mathbb{R}^{n}$ and it satisfies the Lipschitz condition with a non-negative real matrix $K = (k_{ij})_{i,j=1}^{n}$, they construct a sequence of functions $\{x_{m}(t, x_{0})\}$ given by

$$x_m(t,x_0) := x_0 + \frac{1}{\Gamma(p)} \left[\int_0^t (t-s)^{p-1} f(s,x_{m-1}(s,x_0)) ds - \left(\frac{t}{T}\right)^p \int_0^T (T-s)^{p-1} f(s,x_{m-1}(s,x_0)) ds \right],$$
(3)

that satisfies both the differential equation (1) and the boundary conditions (2). Moreover, they prove that the sequence (3) converges uniformly to the exact solution of (1), (2).

This approach was extended in Fečkan and Marynets⁵ to the study of a mixed order FDS

subjected to periodic boundary conditions

2

$$x(0) = x(T), y(0) = y(T),$$
 (5)

with its further application to the fractional-order Duffing equation.¹⁴

The most general case of the periodic FBVP was studied in Fečkan et al,³ where the authors looked at the differential system

$$D_t^p x(t) = f(t, x(t)), \, p \in (m, m+1), \, m \in \mathbb{N},$$
(6)

with periodic boundary conditions

where $t \in [0, T]$, T > 0, $x \in C^m([0, T], D)$, $D \subset \mathbb{R}^n$ is open, $f \in C(G, \mathbb{R}^n)$, $G := [0, T] \times D$. They did not only construct an iterative scheme for approximation of solutions to (6), (7) in the form

$$\begin{aligned} x_{k+1}(t) &= \xi_0 + \sum_{j=1}^m \frac{T^{j-1}}{j!} \left[B_j \left(\frac{t}{T} \right) - B_j \right] \left[-\frac{1}{\Gamma(p-j+1)} \int_0^T (T-s)^{p-j} f(s, x_k(s)) ds \\ &+ \frac{(p-m)T^{m-j+1}}{\Gamma(p-j+2)} \int_0^T (T-s)^{p-m-1} f(s, x_k(s)) ds \right] \\ &+ \frac{1}{\Gamma(p)} \int_0^t (t-s)^{p-1} f(s, x_k(s)) ds - \frac{(p-m)t^p}{T^{p-m}\Gamma(p+1)} \int_0^T (T-s)^{p-m-1} f(s, x_k(s)) ds, \end{aligned}$$
(8)
$$k = 0, 1, \dots, \end{aligned}$$

Even though these results provide us with solutions to the periodic FBVPs, but the conditions we put on the right-hand side of the studied FDEs and on the domain itself are very restrictive.

Motivated by Wang et al,⁷ in this paper, we prove existence and uniqueness of solutions to a higher order FDE in all \mathbb{R} , using techniques of the topological degree (for details, see Gaines and Mawhin¹⁶). The outline of the paper is the following. In Section 2, we formulate the problem setting and prove some auxiliary existence and uniqueness result for a simplified FDE, where the right-hand side does not depend on the unknown function. Section 3 contains the main result that shows existence and uniqueness of solutions of the studied problem.

2 | HIGHER ORDER PERIODIC FBVP

2.1 | Problem setting

Consider the Caputo-type fractional differential equation

$${}_{0}^{C}D_{t}^{q}x(t) = f(t, x(t)), q \in (m - 1, m), m \in \mathbb{N},$$
(9)

subjected to periodic boundary conditions of the form

$$x^{(k)}(0) = x^{(k)}(T), k \in \overline{0, m-1},$$
(10)

where $t \in [0, T]$, with $T > 1, x \in C^{(m-1)}([0, T], \mathbb{R}), f \in C([0, T], \mathbb{R})$ and ${}_{0}^{C}D_{t}^{p}$ being the generalized Caputo fractional derivative with lower limit at 0 (see Podbulny¹²).

Assume that X = C([0, T]) is a Banach space with a maximum norm

$$||x|| = \max_{t \in [0,T]} \{ |x(t)|, |x'(t)|, \dots, |x^{(m-1)}(t)| \},\$$

and let us define operators

$$\mathcal{L}$$
: dom $\mathcal{L} \to X$ and \mathcal{N} : $X \to X$.

with

dom
$$\mathcal{L} = \left\{ x \in C^{(m-1)}([0,T]) : {}_{0}^{C} D_{t}^{q} x(t) \in X, x^{(k)}(0) = x^{(k)}(T), k \in \overline{0, m-1} \right\},\$$

as follows:

$$\mathcal{L}x := {}_0^c D_t^q x \text{ and } \mathcal{N}(x)(t) := f(t, x(t)).$$

Then, periodic FBVP (9), (10) can be rewritten in an operator form

$$\mathcal{L}x = \mathcal{N}(x), x \in \mathrm{dom}\mathcal{L}.$$

2.2 | Auxiliary results and explicit solution

Consider now an auxiliary periodic FBVP:

$${}_{0}^{C}D_{t}^{q}x(t) = z(t), q \in (m-1, m), m \in \mathbb{N},$$
(11)

$$x^{(k)}(0) = x^{(k)}(T), k \in \overline{0, m-1},$$
(10)

with the right-hand side being independent of x(t).

For the problem (11), (10), the following result holds.

Theorem 1. The mapping \mathcal{L} : dom $\mathcal{L} \subset X$ is a Fredholm operator of index zero. Furthermore,

$$Im\mathcal{L} = \left\{ z \in X : \int_{0}^{T} (T-s)^{q-m} z(s) ds = 0 \right\},$$
(12)

 $\ker \mathcal{L} = \{ constant functions \}.$

So if a function $z \in Im\mathcal{L}$, that is, it satisfies a relation:

$$\int_{0}^{T} (T-s)^{q-m} z(s) ds = 0,$$
(13)

then a unique solution $x : [0, T] \rightarrow X$ of the FBVP (11), (10) with (13) is given by

$$x(t) = I^{q} z(t) - \sum_{k=0}^{m-1} b_{k} t^{k},$$
(14)

where

$$b_0 = \frac{\Gamma(q-m+2)}{T^{q-m+1}} I^{2q-m+1} z(T) - (q-m+1) \sum_{k=1}^{m-1} B(q-m+1,k+1) b_k T^k,$$
(15)

$$b_{j+1} = \frac{1}{(j+1)!T} \left[I^{q-j} z(T) - \sum_{k=j+2}^{m-1} \frac{k!}{(k-j)!} b_k T^{k-j} \right], j = \overline{0, m-3},$$
(16)

$$b_{m-1} = \frac{1}{(m-1)!T} I^{q-(m-2)} z(T), \tag{17}$$

with $B(q - m + 1, k + 1) = \frac{\Gamma(q - m + 1)\Gamma(k + 1)}{\Gamma(q - m + k + 2)}$ being the Beta function¹² and $I^p y(t)$ is a Riemann–Liouville fractional integral operator of order p, defined by¹²

$$I^{p}y(t) := \frac{1}{\Gamma(p)} \int_{0}^{t} (t-s)^{p-1}y(s)ds$$

Proof. By Lemma 2.2 in Zhang⁹ for q > 0, the general solution of the homogeneous FDE

$${}_{0}^{c}D_{t}^{q}u(t)=0$$

is given by

$$u(t) = b_0 + b_1 t + \dots + b_{m-1} t^{m-1}, m = [q] + 1,$$

where b_i , $\overline{0, m-1}$ are real constants.

Moreover, using the result of Lemma 2.3 from Zhang,⁹ we deduce that the general solution of the perturbed equation (11) has the form (14), where coefficients b_k , $k = \overline{0, m-1}$ to be defined.

From the (m - 1)th derivative of solution (14), we obtain

$$x^{(m-1)}(t) = I^{q-m+1}z(t) - (m-1)!b_{m-1}$$

Thus, in order for the boundary condition

$$x^{(m-1)}(0) - x^{(m-1)}(T) = 0$$

to hold, function z(t) in the right-hand side of the FDS (11) should satisfy the relation:

$$\int_{0}^{T} (T-s)^{q-m} z(s) ds = 0.$$
 (18)

On the other hand, assume $z \in X$ satisfying (18), and let

$$x(t) = I^q z(t) - \sum_{k=0}^{m-1} b_k t^k.$$

It is easy to prove that x(t), defined by (14), satisfies the rest of the periodic boundary conditions (10), or that $x \in \text{dom }\mathcal{L}$. Indeed, substitution of (14) into the first boundary condition in (10) (for k = 0) gives:

$$x(0) - x(T) = -b_0 - I^q z(T) + b_0 + b_1 T + \sum_{k=2}^{m-1} b_k T^k = 0,$$

$$b_1 = \frac{1}{T} \left[I^q z(T) - \sum_{k=2}^{m-1} b_k T^k \right].$$
(19)

For k = 1, we get

$$x'(0) - x'(T) = -b_1 - I^{q-1}z(T) + b_1 + 2b_2T + \sum_{k=3}^{m-1} kb_kT^{k-1} = 0,$$

$$b_2 = \frac{1}{2!T} \left[I^{q-1}z(T) - \sum_{k=3}^{m-1} kb_kT^{k-1} \right].$$
(20)

For $2 \le j \le m - 3$, we have

$$x^{(j)}(0) - x^{(j)}(T) = -I^{q-j}z(T) + (j+1)!Tb_{j+1} + \sum_{k=j+2}^{m-1} \frac{k!}{(k-j)!} b_k T^{k-j} = 0,$$

$$b_{j+1} = \frac{1}{(j+1)!T} \left[I^{q-j}z(T) - \sum_{k=j+2}^{m-1} \frac{k!}{(k-j)!} b_k T^{k-j} \right].$$
(21)

For k = m - 2,

$$x^{(m-2)}(0) - x^{(m-2)}(T) = -I^{q-(m-2)}z(T) + (m-1)!b_{m-1}T = 0,$$

$$b_{m-1} = \frac{1}{(m-1)!T}I^{q-(m-2)}z(T).$$
(22)

Thus, we conclude that function x(t), defined by the relation (14), satisfies periodic boundary conditions (10), where parameters b_k , $k = \overline{1, m-1}$ are in the form (16), (17). In addition, it follows that

$$\operatorname{Im}\mathcal{L} = \left\{ z \in X : \int_{0}^{T} (T-s)^{q-m} z(s) ds = 0 \right\}.$$
 (23)

Consider now two linear operators \mathcal{P} : $X \to X$ and \mathcal{Q} : $X \to X$ defined by

~

$$\mathcal{P}x(t) = (q - m + 1) \int_{0}^{T} (T - s)^{q - m} x(s) ds, \ t \in [0, T],$$
(24)

5

WILEY

FEČKAN ET AL.

and

6

$$Qz(t) = (q - m + 1) \int_{0}^{T} (T - s)^{q - m} z(s) ds, t \in [0, T].$$
(25)

For $x \in X$, we get

–WILEY-

$$\mathcal{P}(\mathcal{P}x) = \mathcal{P}\left[(q-m+1)\int_{0}^{T} (T-s)^{q-m}x(s)ds\right] = (q-m+1)\int_{0}^{T} (T-s)^{q-m}x(s)ds = \mathcal{P}x$$

Thus, $\mathcal{P}^2 = \mathcal{P}$. Similarly, we obtain that $\mathcal{Q}^2 = \mathcal{Q}$. Note that $\text{Im}\mathcal{P} = \ker \mathcal{L}$ and $\ker \mathcal{Q} = \text{Im}\mathcal{L}$. From the relation

 $\operatorname{ind}\mathcal{L} = \operatorname{dim} \operatorname{ker} \mathcal{L} - \operatorname{codim} \operatorname{ker} \mathcal{L} = 0,$

it follows that
$$\mathcal{L}$$
 is the Fredhold operator of index zero

Since $x \in \ker \mathcal{P}$, that is,

 $(q-m+1)\int_{0}^{T} (T-s)^{q-m}x(s)ds = 0,$ (26)

we deduce that

$$\int_{0}^{T} (T-s)^{q-m} \left[I^{q} z(s) - b_{0} - \sum_{k=1}^{m-1} b_{k} s^{k} \right] ds = 0,$$
(27)

and thus,

$$b_0 = \frac{\Gamma(q-m+2)}{T^{q-m+1}} I^{2q-m+1} z(T) - (q-m+1) \sum_{k=1}^{m-1} B(q-m+1,k+1) b_k T^k,$$
(28)

where b_k , $k = \overline{1, m - 1}$ are defined by (16), (17).

Substituting b_0 into (14), we obtain the unique solution to the FBVP (11), (10).

Example. To verify the result of Theorem 1, consider a periodic FBVP of the form:

$${}_{0}^{C}D_{t}^{2.5}x(t) = -3t + 2 \ (:= z(t)), \ t \in [0, 1],$$
⁽²⁹⁾

$$x(0) = x(1), \ x'(0) = x'(1), \ x''(0) = x''(1),$$
(30)

where m = 3.

It is easy to check that function z(t) satisfies condition (13), and thus, solution x(t) of (29), (30) can be written in the form:

$$x(t) = I^{2.5} z(t) - b_0 - b_1 t - b_2 t^2,$$

where parameters b_i , $i = \overline{0, 2}$ to be calculated following the process presented in the proof of Theorem 1.

Using mathematical software Maple 2022, we find that

$$b_0 = \frac{\pi}{2} + \frac{352}{105}, \ b_1 = \frac{8}{105\sqrt{\pi}}, \ b_2 = \frac{8}{15\sqrt{\pi}}$$

and thus, solution x(t) of the periodic FBVP (29), (30) can be written as

$$x(t) = \frac{-96t^{3.5} + 224t^{2.5} - 112t^2 - 16t - 704\pi^{0.5} - 105\pi^{1.5}}{210\sqrt{\pi}}.$$
(31)

Direct substitution of (31) into (29), (30) shows that x(t) indeed satisfies both. As an additional verification, we use the fact that integration of the FDE (29) leads to a solution of the form:

$$x(t) = x(0) + tx'(0) + \frac{t^2}{2}x''(0) + I^{2.5}z(t).$$
(32)

Substituting (31) into (32) and comparing its left- and right-hand sides, we conclude that they coincide.

Thus, we have demonstrated on a concrete example of a periodic FBVP that under conditions of Theorem 1, solution to the problem indeed has the form (14) with coefficients b_i , $i = \overline{0, m-1}$ being calculated according to formulas (15)–(17).

3 | EXISTENCE RESULT

Consider now a projection \mathcal{P} : $X \to X$ defined by

$$\mathcal{P}x(t) = (q - m + 1) \int_{0}^{T} (T - s)^{q - m} x(s) ds, t \in [0, T].$$
(33)

Let us set $Q = \mathcal{I} - \mathcal{P}$ and note that ker $Q = \ker \mathcal{L}$ and ker $\mathcal{P} = \operatorname{Im}\mathcal{L}$. Additionally, we denote by \mathcal{L}^{-1} : ker $\mathcal{P} \to \ker \mathcal{P} \cap \operatorname{dom}\mathcal{L} \subset X$ an inverse operator given by (14).

The following existence result holds.

Theorem 2. Assume that there exist positive constants M_1 and k_0 such that $|f(t,x)| \le M_1$, for $t \in [0,T]$, $x \in \mathbb{R}$, and

$$either$$

$$\int_{0}^{T} (T-s)^{q-m} f(s,x) ds > 0, \ \forall x \ge k_{0}, \ \int_{0}^{T} (T-s)^{q-m} f(s,x) ds < 0, \ \forall x \le -k_{0},$$
or
$$\int_{0}^{T} (T-s)^{q-m} f(s,x) ds < 0, \ \forall x \ge k_{0}, \ \int_{0}^{T} (T-s)^{q-m} f(s,x) ds > 0, \ \forall x \le -k_{0}.$$
(34)

Then, the problem (9), (10) has at least one solution.

Proof. From Theorem 1, it follows that function x(t) is a solution of the periodic FBVP (9), (10) if and only if $\mathcal{PN}(x) = 0$ and then $Qx = \mathcal{L}^{-1}Q\mathcal{N}(x)$, which is equivalent to an equation

$$\mathcal{T}(x) = \mathcal{P}\mathcal{N}(x) + Qx - \mathcal{L}^{-1}Q\mathcal{N}(x) = 0.$$
(35)

Note that $Qx - \mathcal{L}^{-1}Q\mathcal{N}(x) \in \ker \mathcal{P}$.

Next, we will show that the operator $\mathcal{T} : X \to X$ is completely continuous. Let $B \subset X$ be a bounded set. By the assumption that $|f(t, x(t))| \le M_1$, for $x \in B$, we deduce

$$\begin{aligned} |\mathcal{T}x(t)| &\leq \frac{1}{\Gamma(q)} \int_{0}^{t} (t-s)^{q-1} |f(s,x(s))| ds + |b_0| + \sum_{k=1}^{m-1} |b_k| T^k \\ &\leq M_1 T^q \left[\frac{1}{\Gamma(q+1)} + \frac{\Gamma(q-m+2)}{\Gamma(2q-m+2)} + \sum_{k=1}^{m-1} \left\{ \frac{\Gamma(q-m+2)}{\Gamma(q-m+k+2)\Gamma(q-k)} + \frac{1}{k!\Gamma(q-k)} \right\} \right] := M_2. \end{aligned}$$
(36)

Indeed, calculations show that the following estimates hold:

$$\begin{split} \frac{1}{\Gamma(q)} \int_{0}^{t} (t-s)^{q-1} |f(s,x(s))| ds &\leq \frac{M_1}{\Gamma(q)} \int_{0}^{t} (t-s)^{q-1} ds = \frac{M_1 t^q}{\Gamma(q+1)} \leq \frac{M_1 T^q}{\Gamma(q+1)} \\ |b_{m-1}| T^{m-1} &\leq \frac{T^{m-2}}{(m-1)! \Gamma(q-m+2)} \int_{0}^{T} (T-s)^{q-m+1} |f(s,x(s))| ds \\ &\leq \frac{M_1 T^{m-2}}{(m-1)! \Gamma(q-m+2)} \int_{0}^{T} (T-s)^{q-m+1} ds = \frac{M_1 T^q}{(m-1)! \Gamma(q-m+3)}, \end{split}$$

and thus,

8

ΊΙΕV

$$\begin{split} |b_{m-1}| &\leq \frac{M_1 T^{q-m+1}}{(m-1)!\Gamma(q-m+1)} \cdot \frac{1}{(q-m+2)(q-m+1)} < \frac{M_1 T^{q-m+1}}{(m-1)!\Gamma(q-m+1)}, \\ |b_{m-2}| &\leq \frac{M_1 T^{q-m+2}}{(m-2)!\Gamma(q-m+2)} \left[\frac{1}{(q-m+3)(q-m+2)} + \frac{1}{2!(q-m+2)} \right] < \frac{M_1 T^{q-m+2}}{(m-2)!\Gamma(q-m+2)}, \\ |b_{m-3}| &\leq \frac{M_1 T^{q-m+3}}{(m-3)!\Gamma(q-m+3)} \left[\frac{1}{(q-m+3)(q-m+4)} + \frac{1}{2!} \left\{ \frac{1}{(q-m+3)} + \frac{1}{2!} \right\} + \frac{1}{3!} \right] < \frac{M_1 T^{q-m+3}}{(m-3)!\Gamma(q-m+3)} \end{split}$$

Continuing computations further, we obtain that

$$\begin{split} |b_j| &\leq \frac{M_1 T^{q-j}}{j! \Gamma(q-j)}, j = \overline{1, m-1}, \\ |b_0| &\leq \frac{\Gamma(q-m+2)M_1 T^q}{\Gamma(2q-m+2)} + \sum_{k=1}^{m-1} \frac{\Gamma(q-m+2)M_1 T^q}{\Gamma(q-m+k+2)\Gamma(q-k)} \end{split}$$

Furthermore, calculations show that the inequality holds:

$$\begin{aligned} |(\mathcal{T}x)'(t)| &\leq \frac{1}{\Gamma(q-1)} \int_{0}^{t} (t-s)^{q-2} |f(s,x(s))| ds + \sum_{k=1}^{m-1} |kb_{k}| T^{k-1} \\ &\leq M_{1} T^{q-1} \left[\frac{1}{\Gamma(q)} + \sum_{k=1}^{m-1} \frac{1}{(k-1)!\Gamma(q-k)} \right] := M_{3}. \end{aligned}$$
(37)

Hence, for $t_1, t_2 \in [0, T]$, we conclude that

$$|(\mathcal{T}x)(t_2) - (\mathcal{T}x)(t_1)| \le \int_{t_1}^{t_2} |(\mathcal{T}x)'(s)| ds \le M_3(t_2 - t_1).$$
(38)

This proves that \mathcal{T} is equicontinuous on [0, T]. Hence, by Arzela–Ascoli theorem, the operator $\mathcal{T} : X \to X$ is completely continuous.

Now, we need to show that there exists at least one solution $x \in C[0, T]$ satisfying (35). For this purpose, we take a homotopy

$$\mathcal{T}_{\lambda}(x) = \mathcal{PN}(x) + Qx - \lambda \mathcal{L}^{-1} \mathcal{QN}(x), \ \lambda \in [0, 1],$$
(39)

and consider the set

$$\Omega = \{ x \in X : \|\mathcal{P}x\| \le k_1, \|\mathcal{Q}x\| \le k_2 \}$$

for

$$k_1 = k_0 + k_2, \ k_2 = \|\mathcal{L}^{-1}\| \|\mathcal{Q}\| M_1 + 1.$$

10991476, 0, Downloaded from https://onlinelibrary.wiley.com/doi/10.1002/nma.9097 by Tu Defft, Wiley Online Library on [13/02/2023]. See the Terms and Conditions (https://onlinelibrary.wiley.com/terms-and-conditions) on Wiley Online Library for rules of use; OA articles are governed by the applicable Creative Commons License

Then, we claim

 $\mathcal{T}_{\lambda}(x) \neq 0, \forall x \in \partial \Omega.$

Indeed, if $\mathcal{T}_{\lambda}(x) = 0$ for $||Qx|| = k_2$, then

$$\|\mathcal{L}^{-1}\|Q\|M_1 + 1 = k_2 = \|Qx\| = \|\lambda \mathcal{L}^{-1}Q\mathcal{N}(x)\| \le \|\mathcal{L}^{-1}\|\|Q\|M_1$$

which is a contradiction. If $\mathcal{T}_{\lambda}(x) = 0$ for $||\mathcal{P}x|| = k_1$, then

$$|x(t)| \ge |\mathcal{P}x| - ||\mathcal{Q}x|| \ge k_1 - k_2 = k_0$$

and

$$0 = \int_{0}^{T} (T - s)^{q - m} f(s, x(s)) ds \neq 0,$$

by (34), which is again a contradiction.

Thus, we have

$$\deg(\mathcal{T}, \Omega, 0) = \deg(\mathcal{T}_0, \Omega, 0) = \deg(\mathcal{PN} + \mathcal{Q}, \Omega, 0)$$
$$= \deg(\mathcal{PN}, \Omega \cap \ker \mathcal{N}, 0) = \deg\left(\int_0^T (T - s)^{q - m} f(s, x) ds, (-k_0, k_0), 0\right) = \pm 1 \neq 0.$$

This means that the operator \mathcal{T} has at least one zero point in Ω , which implies that (9), (10) has at least one solution.

We present an example. Consider the Caputo-type fractional differential equation

$${}_{0}^{C}D_{t}^{q}x(t) = \mu \tanh x(t) + \nu \cos t, \ q \in (m-1,m), \ m \in \mathbb{N},$$
(41)

for $\mu, \nu \in \mathbb{R} \setminus \{0\}$ and subjected to periodic boundary conditions

$$x^{(k)}(0) = x^{(k)}(2\pi), k \in \overline{0, m-1}.$$
(42)

We set $\alpha = q - m \in (0, 1)$. Then, $T = 2\pi$, $f(x, t) = \mu \tanh x + v \cos t$, and hence,

$$\int_{0}^{2\pi} (2\pi - s)^{\alpha} f(s, x) ds = \int_{0}^{2\pi} (2\pi - s)^{\alpha} (\mu \tanh x + \nu \cos s) ds = \mu \tanh x \int_{0}^{2\pi} (2\pi - s)^{\alpha} ds + \nu \int_{0}^{2\pi} (2\pi - s)^{\alpha} \cos s) ds$$
$$= \mu \frac{(2\pi)^{\alpha+1}}{\alpha+1} \tanh x + \nu \int_{0}^{2\pi} s^{\alpha} \cos s ds \rightarrow_{x \to \pm \infty} \pm \mu \frac{(2\pi)^{\alpha+1}}{\alpha+1} + \nu \int_{0}^{2\pi} s^{\alpha} \cos s ds.$$

Thus, condition (34) is satisfied if it holds

 $-\mu^2 \frac{(2\pi)^{2(\alpha+1)}}{(\alpha+1)^2} + \nu^2 \left(\int_{0}^{2\pi} s^{\alpha} \cos s ds\right)^2 < 0,$

which is equivalent to

$$\frac{\alpha+1}{(2\pi)^{\alpha+1}} \left| \int_{0}^{2\pi} s^{\alpha} \cos s ds \right| = \left| {}_{1}F_{2} \left(1; \frac{a}{2} + 1, \frac{a}{2} + \frac{3}{2}; -\pi^{2} \right) \right| < \frac{|\mu|}{|\nu|},$$
(43)

where $_1F_2$ is a generalized hypergeometric function. By applying Theorem 2, we arrive at the following result.

(40)

WII FY



FIGURE 1 Graph of (43). [Colour figure can be viewed at wileyonlinelibrary.com]

Theorem 3. If condition (43) holds, then (41) with (42) has a solution.

Using the expansion

$$\int_{0}^{2\pi} s^{\alpha} \cos s ds = \sum_{k=0}^{\infty} (-1)^{k} \frac{(2\pi)^{2k+\alpha+1}}{(2k)!(2k+\alpha+1)},$$

we can approximately compute (43) with the graph on Figure 1.

4 | SYSTEMS OF FDE

The result of Theorem 2 can be generalized to the case of systems of FDEs. So we consider the case $f \in C([0, T], \mathbb{R}^n)$ for $n \ge 2$. Then, conditions (34) need to be modified by evaluating the quantity |x| instead of x itself. To be more concrete, we have the following results.

Theorem 4. Assume that there exist positive constants M_1 and k_0 such that $|f(t,x)| \le M_1$, for $t \in [0, T]$, $x \in \mathbb{R}^n$ such that

$$\int_{0}^{T} (T-s)^{q-m} f(s,x) ds \neq 0, \, \forall |x| \ge k_0,$$
(44)

and

$$\deg\left(\int_{0}^{T} (T-s)^{q-m} f(s,x) ds, B(k_0), 0\right) \neq 0,$$
(45)

for the ball $B(k_0) = \{x \in \mathbb{R}^n : |x| \le k_0\}$. Then, the problem (9), (10) has at least one solution.

Proof. The result follows directly from the proof of Theorem 2. **Theorem 5.** Assume there is a scalar product (\cdot, \cdot) : $\mathbb{R}^n \times \mathbb{R}^n \to \mathbb{R}$ and a positive constant k_0 such that

$$\int_{0}^{T} (T-s)^{q-m} (f(s,x)ds,x) \neq 0, \, \forall |x| \ge k_0.$$
(46)

Then, the problem (9), (10) has at least one solution.

Proof. We apply Theorem 4. Clearly, (46) implies (44). Since the set $\{x \in \mathbb{R}^n : |x| \ge k_0\}$ is connecting, we have the following two possibilities:

1. $\int_0^T (T-s)^{q-m} (f(s,x)ds,x) > 0, \forall |x| \ge k_0$. Then, we derive

$$\left(\lambda \int_{0}^{T} (T-s)^{q-m} f(s,x) ds + (1-\lambda)x\right) = \lambda \int_{0}^{T} (T-s)^{q-m} (f(s,x) ds,x) + (1-\lambda)(x,x) > 0,$$

for $\lambda \in [0, 1]$ and $|x| \ge k_0$. This implies

$$\deg\left(\int_{0}^{T} (T-s)^{q-m} f(s,x) ds, B(k_0), 0\right) = \deg\left(I, B(k_0), 0\right) = 1 \neq 0.$$

Hence, (45) holds.

2. $\int_0^T (T-s)^{q-m} (f(s,x)ds,x) < 0, \forall |x| \ge k_0.$ Then, we derive

$$\left(\lambda \int_{0}^{T} (T-s)^{q-m} f(s,x) ds - (1-\lambda)x\right) = \lambda \int_{0}^{T} (T-s)^{q-m} (f(s,x) ds,x) - (1-\lambda)(x,x) < 0,$$

for $\lambda \in [0, 1]$ and $|x| \ge k_0$. This implies

$$\deg\left(\int_{0}^{T} (T-s)^{q-m} f(s,x) ds, B(k_0), 0\right) = \deg\left(-I, B(k_0), 0\right) = (-1)^n \neq 0.$$

Hence, (45) holds. The proof is finished.

ACKNOWLEDGEMENTS

The authors are grateful to the reviews for their valuable comments that helped to improve the paper.

CONFLICT OF INTEREST STATEMENT

This work does not have any conflicts of interest.

ORCID

Michal Fečkan https://orcid.org/0000-0002-7385-6737 Kateryna Marynets https://orcid.org/0000-0002-0043-6336 JinRong Wang https://orcid.org/0000-0002-6642-1946

REFERENCES

- 1. Farkas M. Periodic Motions. Springer-Verlag; 1994.
- 2. Hu L, Zhang S. Existence of positive solutions of a periodic boundary value problems for nonlinear fractional differential equations at resonance. *J Frac Calc Appl.* 2017;8(2):19-31.
- Fečkan M, Marynets K, Wang JR. Periodic boundary value problems for higher order fractional differential systems. *Math Meth Appl Sci.* 2019;42:3616-3632.
- 4. Fečkan M, Marynets K. Approximation approach to periodic BVP for mixed fractional differential systems. *J Comp Appl Math.* 2018;339:208-217.
- 5. Fečkan M, Marynets K. Approximation approach to periodic BVP for fractional differential systems. *Eur Phys Jour: Special Topics*. 2017;226:3681-3692.
- 6. Hu Z, Liu W, Rui W. Periodic boundary value problem for fractional differential equation. Int J Math. 2012;23:1250100.
- 7. Wang JR, Ibrahim AG, Fečkan M. Differential inclusions of arbitrary fractional order with anti-periodic conditions in Banach space. *Electron J Qual Theory Differ Equ.* 2016;34:1-22.

¹²WILEY

- 8. Ahmad B, Nieto JJ. Existence of solutions for anti-periodic boundary value problems involving fractional differential equations via Leray-Schauder degree theory. *Topol Meth Nonl Anal.* 2010;35:235-304.
- 9. Zhang S. Positive solutions for boundary value problems of nonlinear fractional differential equations. *Electron J Differ Equ.* 2006;2006:1-12.
- 10. Marynets K. Solvability analysis of a special type fractional differential system. Com Appl Math. 2020;39(1):1-13.
- 11. Kilbas A, Srivastava HM, Trujillo J. Theory and Applications of Fractional Differential Equations. Elsevier; 2006.
- 12. Podbulny I. Fractional Differential Equations, 1st ed. Academic Press; 1999.
- 13. Zhou Y. Basic Theory of Fractional Differential Equations. World Scientific; 2014.
- 14. Li Z, Chen D, Zhu J, Liu Y. Nonlinear dynamics of fractional order Duffing system. Chaos Solitons Fractals. 2015;81:111-116.
- 15. Sebah P, Gourdon X. Introduction to Bernoulli's numbers; 2012.
- 16. Gaines RE, Mawhin JL. Coincidence Degree and Nonlinear Differential Equations. Springer; 1977.

How to cite this article: Fečkan M, Marynets K, Wang J. Existence of solutions to the generalized periodic fractional boundary value problem. *Math Meth Appl Sci.* 2023;1-12. doi:10.1002/mma.9097