

## Scenarios for Controls of River Response to Climate Change in the Lower Rhine River

Ylla Arbos, Claudia; Blom, Astrid; Schielen, Ralph M.J.

**Publication date**

2021

**Document Version**

Final published version

**Citation (APA)**

Ylla Arbos, C., Blom, A., & Schielen, R. M. J. (2021). *Scenarios for Controls of River Response to Climate Change in the Lower Rhine River*. 70-71. Abstract from NCR DAYS 2021. <https://kbase.ncr-web.org/outputs/rivers-in-an-uncertain-future/>

**Important note**

To cite this publication, please use the final published version (if applicable). Please check the document version above.

**Copyright**

Other than for strictly personal use, it is not permitted to download, forward or distribute the text or part of it, without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license such as Creative Commons.

**Takedown policy**

Please contact us and provide details if you believe this document breaches copyrights. We will remove access to the work immediately and investigate your claim.

***Green Open Access added to TU Delft Institutional Repository***

***'You share, we take care!' - Taverne project***

**<https://www.openaccess.nl/en/you-share-we-take-care>**

Otherwise as indicated in the copyright section: the publisher is the copyright holder of this work and the author uses the Dutch legislation to make this work public.

Book of abstracts

# Rivers in an uncertain future

NCR days 2021 | Enschede, February 11-12



Jord J. Warmink, Anouk Bomers,  
Vasileios Kitsikoudis, R. Pepijn van  
Denderen & Fredrik Huthoff (eds.)

NCR publication: 46-2021

Netherlands  
Centre for  
River studies **NCR**

# NCR Days 2021

*Rivers in an uncertain future*

Jord J. Warmink, Anouk Bomers, Vasileios Kitsikoudis, R. Pepijn  
van Denderen & Fredrik Huthoff (eds.)

Organising partner:

## UNIVERSITY OF TWENTE.

Co-sponsored by:



### Organising partner

University of Twente  
P.O. 217  
7500 AE Enschede  
The Netherlands

telephone: +31 53 489 35 46  
e-mail: [secretariat-mfs-et@utwente.nl](mailto:secretariat-mfs-et@utwente.nl)  
www: <https://www.utwente.nl>

### Contact NCR

dr. ir. K.D. Berends (programme Secretary)  
Netherlands Centre for River Studies  
c/o Deltares  
P.O. 177  
2600 MH Delft  
The Netherlands

telephone: +31 6 21 28 74 61  
e-mail: [secretary@ncr-web.org](mailto:secretary@ncr-web.org)  
www: <https://www.ncr-web.org>

**Cite as:** Warmink, J.J., Bomers, A., Kitsikoudis, V., van Denderen, R.P., Huthoff, F. (2021), *Rivers in an uncertain future: NCR Days proceedings*. Netherlands Centre for River Studies, publication 46-2021

**Photo credits cover:** IJssel river from Bureau Beeldtaal Filmmakers (2018)

Copyright © 2021 Netherlands Centre for River Studies

All rights reserved. No parts of this document may be reproduced in any form by print, photo print, photo copy, microfilm or any other means, without permission of the publisher: Netherlands Centre for River Studies.

# Scenarios for Controls of River Response to Climate Change in the Lower Rhine River

Clàudia Ylla Arbós<sup>a,\*</sup>, Astrid Blom<sup>a</sup>, Ralph M.J. Schielen<sup>a,b</sup>

<sup>a</sup>*Delft University of Technology, Department of Hydraulic Engineering, Faculty of Civil Engineering and Geosciences, PO Box 5048, 2600 GA Delft, the Netherlands*

<sup>b</sup>*Ministry of Infrastructure and Water Management - DG Rijkswaterstaat, Utrecht, the Netherlands*

**Keywords** — Climate Change, River Controls, Lower Rhine River

## Introduction

The majority of the world's large rivers are heavily engineered. In such heavily engineered rivers, channel response (i.e., changes in bed elevation, channel slope, and bed surface grain size) is predominantly determined by human intervention [e.g., Surian and Rinaldi (2003); Ylla Arbós et al. (2020)]. The relative influence of climate on channel response may, however, increase in the upcoming decades, as climate change alters the river controls. Specifically, climate change affects (1) the characteristics of the flow rate, through changes in precipitation patterns; (2) the downstream base level, through sea level rise, or lake base level drop; and (3) the sediment flux, due to changes in water discharge.

It is increasingly necessary to anticipate future channel response to climate-related changes in the river controls. This can be done using numerical models, by changing their boundary conditions. A required step is determining climate change scenarios for the river controls and translating them into suitable boundary conditions for numerical models, taking into account different sources of uncertainty.

Here we consider the Lower Rhine River, from Bonn (Germany) to Gorinchem (Netherlands). We discuss the projected changes of the hydrodynamic river controls over the 21<sup>st</sup> century, and how they can be transformed into suitable boundary conditions for a schematized model.

## Changes in water discharge

Several studies have attempted to predict water discharge in the Rhine basin until 2100 using a model chain [e.g., Gorgen (2010); Hegnauer (2017)]. First, the different emission scenarios or representative concentration pathways (RCP's) defined by IPCC are used as input to climate models. The output is then used to obtain future precipitation time series. Finally, the precipitation time series are input

to a hydrological model that transforms daily precipitation into daily river discharge. Numerous models and techniques are used in this process, and the results are sensitive to these choices, in particular to that of the climate model (Gorgen, 2010; Hegnauer, 2017). In general terms, the different studies predict similar future trends in water discharge along the Rhine basin (Hegnauer, 2017). In upstream, snowmelt-dominated locations (e.g., Basel), higher temperatures lead to earlier snowmelt, which slightly shifts the annual discharge peak to earlier in the year, the magnitude of the peak not changing significantly. In the downstream, rainfall-controlled tributaries (e.g. Trier, Raunheim), higher peak flow rates and lower base flow rates are expected, without shifts in the timing of the peaks. In the Lower Rhine River, a combined behavior is expected, with a rainfall-peak in the winter, a smaller snowmelt-peak in summer, and higher peak flow rates and lower base flow rates. At Bonn, the winter mean monthly discharge may increase 15-50%, while summer mean monthly discharge may decrease 0-40% (Gorgen, 2010; Hegnauer, 2017).

A river tends to equilibrium channel characteristics, eventually attained if the controls do not change, or do so at low rates (Blom et al., 2017). The equilibrium bed profile can be decomposed in a mean bed profile, and fluctuations about it. The mean bed profile depends on the combination magnitude-frequency of flow events (the flow duration curve), while the order of flow events determines the fluctuations (Arkesteijn et al., 2019). Channel response is inseparable from discharge variability, and more information on discharge variability than mean monthly discharge is needed as an upstream flow boundary condition.

Gorgen (2010) provides an ensemble of synthetic daily discharge time series covering the period 1951-2100, and Hegnauer (2017) provides 50-year synthetic daily discharge time series representative of the period 1951-2006, 2050, and 2085. We propose to use these data to create simplified hydrographs, which sufficiently preserve discharge variability, while ensuring that the only difference between sce-

\*Corresponding author

Email address: c.yllaarbos@tudelft.nl (Clàudia Ylla Arbós)

URL: www.tudelft.nl (Clàudia Ylla Arbós)



narios is the climate signal, facilitating comparison between scenarios. A base-case, cycled 10-year hydrograph with similar statistical properties as the long-term historic time series, can be modified for each climate scenario, such that the future hydrographs have the same statistical properties as the synthetic hydrographs provided by Gorgen (2010); Hegnauer (2017), but keep the the order of flow events unchanged. From the synthetic hydrographs, the probability density function of water discharge for each scenario  $S_i$  can be obtained, providing the discharge associated to a certain probability of occurrence  $p$ , that is,  $Q_{S_i}(p)$  (Figure 1a). The same exercise can be done with the base-case hydrograph, so as to obtain  $Q_0(p)$ . Each flow event in the base-case hydrograph can then be related to a probability of occurrence  $p$ , and further multiplied by the factor  $F_{S_i}(p) = Q_{S_i}(p) / Q_0(p)$  (Figure 1b). The effect of the hydrograph cycled period can then be assessed. Further research is needed to validate this method.




Figure 1: (a) Statistics of water discharge in the base case, and scenario  $S_i$ ; (b) simplified hydrograph in the base case and in scenario  $S_i$ .

### Changes in sea level

Climate scenarios foresee an increase of sea level between 0.2 and 3 m by the end of the century (Haasnoot et al., 2018). In all scenarios, sea level rise accelerates with time (up to 7 cm/a) when compared to the current rates of 0.2 cm/a. The increase in sea level has to be translated into the corresponding increase of water surface elevation at Gorinchem, the downstream boundary of our domain of interest. As an indication, an increase of sea level of 3 m at Hoek van Holland, would result into a water level increase of 2.75 m at Gorinchem, considering a water discharge of 2200 m<sup>3</sup>/s, and based on hydrodynamic computations only (Haasnoot et al., 2018).

An increase in base level is expected to trigger a morphodynamic response towards a new equilibrium state, which consists of an increase of bed level equal to the total rise of sea level.

Due to the sand deficit in the Lower Rhine-Meuse Delta (Cox et al., 2020), bed level cannot keep pace with sea level rise. Further analysis will provide insight on how the water level at Gorinchem will be affected by sea level rise.

### Discussion and conclusions

Climate change affects the boundary conditions of the Lower Rhine River, which may affect future channel response. We have discussed foreseen changes in the hydrodynamic river controls. In addition, scenarios for changes in the sediment flux are required, based on changes in the hydrograph, and land-use changes. The high degree of uncertainty related to climate projections calls for a statistical or a scenario approach.

Changes in future intervention policy may have a greater effect on channel response than climate change. Modeling of channel response must consider intervention scenarios. A comparison between climate-triggered response and intervention-triggered response will shed light on the relative importance of natural versus human controls on channel response.

### References

- Arkesteijn, L., Blom, A., Czapiga, M. J., Chavarrías, V., and Labeur, R. J. (2019). The Quasi-Equilibrium Longitudinal Profile in Backwater Reaches of the Engineered Alluvial River: A Space-Marching Method. *JGR:ES*, 124(11):2542–2560.
- Blom, A., Arkesteijn, L., Chavarrías, V., and Viparelli, E. (2017). The equilibrium alluvial river under variable flow and its channel-forming discharge. *JGR:ES*, 122(10):1924–1948.
- Cox, J., Kleinhans, M. G., and Huismans, Y. (2020). An estuary out of equilibrium: The importance of dredging in determining the net sediment flux in the Rhine-Meuse Estuary. *River Flow 2020*, Delft.
- Görgen, K. (2010). Assessment of climate change impacts on discharge in the Rhine River Basin : results of the RheinBlick2050 project.
- Haasnoot, M., Bouwer, L., Diermanse, F., Kwadijk, J., van der Spek, A., (...) Huismans, Y., Sloff, K., and Mosselman, E. (2018). Mogelijke gevolgen van versnelde zeespiegelstijging voor het Deltaprogramma. Een verkenning.
- Hegnauer, M. (2017). Analysis GRADE results for different locations in the Rhine Basin. *Deltares*.
- Surian, N. and Rinaldi, M. (2003). Morphological response to river engineering and management in alluvial channels in Italy. *Geomorphology*, 50(4):307–326.
- Ylla Arbós, C., Blom, A., Viparelli, E., Reenekens, M., Frings, R. M., and Schielen, R.M.J. (2020). River Response to Anthropogenic Modification: Channel Steepening and Gravel Front Fading in an Incising River. *GRL*. doi:10.1029/2020GL091338