

Geodynamics of the Romanian Territory Inferred from a Dense Continuous GNSS Network

Muntean, A.; Ambrosius, B.A.C.; Mocanu, V.; Nastase, Eduard Ilie

DOI

[10.3997/2214-4609.202149BGS92](https://doi.org/10.3997/2214-4609.202149BGS92)

Publication date

2021

Document Version

Final published version

Citation (APA)

Muntean, A., Ambrosius, B. A. C., Mocanu, V., & Nastase, E. I. (2021). *Geodynamics of the Romanian Territory Inferred from a Dense Continuous GNSS Network*. Abstract from 11th Congress of the Balkan Geophysical Society, Bucharest, Romania. <https://doi.org/10.3997/2214-4609.202149BGS92>

Important note

To cite this publication, please use the final published version (if applicable).
Please check the document version above.

Copyright

Other than for strictly personal use, it is not permitted to download, forward or distribute the text or part of it, without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license such as Creative Commons.

Takedown policy

Please contact us and provide details if you believe this document breaches copyrights.
We will remove access to the work immediately and investigate your claim.

Geodynamics of the Romanian Territory Inferred from a Dense Continuous GNSS Network

A. Muntean¹, B. Ambrosius², V. Mocanu³, E. Nastase¹

¹ National Institute for Earth Physics; ² Delft University of Technology, Faculty Aerospace Engineering; ³ University of Bucharest, Department of Geophysics

Summary

No Summary provided

Geodynamics of the Romanian territory inferred from a dense continuous GNSS network

A. Muntean¹, B.A.C Ambrosius², V. Mocanu³, E. Nastase^{1,3}

¹ *National Institute for Earth Physics, Romania*

² *Delft University of Technology, Faculty Aerospace Engineering, Netherlands*

³ *University of Bucharest, Department of Geophysics, Romania*

Abstract

Over the last two decades the Romanian territory has been covered by a dense permanent Global Navigation Satellite System (GNSS) network, presently including more than one hundred stations established and maintained by several Romanian agencies. We refer to these as continuous GNSS (cGNSS) stations. For the study presented here we exclusively use the data of the Global Positioning System (GPS), which is the oldest and most widely-used element of GNSS. At the beginning of this century, we realized the potential of cGNSS for scientific purposes. Applications include monitoring of crustal deformations associated with the Carpathian orogen and the neighbouring platforms in Romania, in correlation with tectonic processes in South-East Europe (Africa-Europe plate interactions), observation of (vertical) crustal motions to study the surface expression of deep earthquakes in the Eastern Carpathians Bending Zone (Vrancea region).

This study presents the results of our analysis of the aggregated cGNSS measurements in Romania. The results cover the period 2000-2020, but focus on the period 2013-2020 when the majority of the stations became operational. We processed the Receiver Independent Exchange Format (RINEX) daily data (sampled at 30-second intervals) of all available stations. For this purpose, we used the GipsyX scientific GPS analysis software, applying the so-called Precise Point Positioning (PPP) strategy. This resulted in time-series of the horizontal and vertical position components of each station in the International Terrestrial Reference Frame 2014 (ITRF14). Subsequently, the time-series were converted to a stable Eurasian tectonic reference plate using the ITRF14 plate rotation model for Eurasia. All data processing in this study is based on this reference frame.

The results of this complex, long-term study indicates that the Romanian territory trends to move slightly southward relative to stable Eurasia, at velocity rates of about 1.0 mm/yr in the north and up to 3.0 mm/yr in the south. This may be an indication of plate-stretching due to slab-rollback of the African plate relative to the Eurasian plate. Until now we did not find an obvious correlation between the deep Vrancea earthquakes and (vertical) surface motions. That is probably because the recent earthquakes were too small.

Acknowledgements

This research was carried out within the NUCLEU project, MULTIRISC Program that is supported by the Ministry of Research and Innovation, project nr. PN19080201. The GipsyX software is licensed to the Department of Geophysics of the University of Bucharest (UNIBUC). Other institutions could apply this software in an ongoing collaboration with UNIBUC. The authors thank the National Research and Development Institute for Marine Geology and Geoecology, National Center for Monitoring and Alarm to Natural Marine Hazards – Euxinus, and the National Agency for Cadaster and Land Registration for providing access to their data.