

# Dynamics of Thioalkalivibrio species in a co-culture under selective pressure of ampicillin

Ahn, Anne Catherine; Schuurmans, J. Merijn; Sorokin, Dimitry; Muyzer, Gerard

DOI 10.1093/femsmc/xtad020

Publication date 2023 **Document Version** Final published version

Published in **FEMS Microbes** 

Citation (APA) Ahn, A. C., Schuurmans, J. M., Sorokin, D., & Muyzer, G. (2023). Dynamics of Thioalkalivibrio species in a co-culture under selective pressure of ampicillin. FEMS Microbes, 4. https://doi.org/10.1093/femsmc/xtad020

### Important note

To cite this publication, please use the final published version (if applicable). Please check the document version above.

Copyright

Other than for strictly personal use, it is not permitted to download, forward or distribute the text or part of it, without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license such as Creative Commons.

Takedown policy

Please contact us and provide details if you believe this document breaches copyrights. We will remove access to the work immediately and investigate your claim.

DOI: 10.1093/femsmc/xtad020 Advance access publication date: 8 November 2023 Research Article – Microbes & Environment

# Dynamics of Thioalkalivibrio species in a co-culture under selective pressure of ampicillin

Anne-Catherine Ahn<sup>1,2</sup>, J. Merijn Schuurmans<sup>1</sup>, Dimitry Sorokin<sup>3,4</sup>, Gerard Muyzer<sup>1,\*</sup>

<sup>1</sup>Microbial Systems Ecology, Department of Freshwater and Marine Ecology, Institute for Biodiversity and Ecosystem Dynamics, University of Amsterdam, 1098 XH, Amsterdam, The Netherlands

<sup>2</sup>Current address: Laboratory of Microbiology, Wageningen University & Research, Stippeneng 4, 6708 WE, Wageningen, The Netherlands

<sup>3</sup>Winogradsky Institute of Microbiology, Federal Research Centre of Biotechnology, Russian Academy of Sciences, 119071, Moscow, Russia

<sup>4</sup>Department of Biotechnology, Delft University of Technology, 2629 HZ, Delft, The Netherlands

\*Corresponding author. Microbial Systems Ecology, Department of Freshwater and Marine Ecology, Institute for Biodiversity and Ecosystem Dynamics, University of Amsterdam, 1098 XH, Amsterdam, The Netherlands. E-mail: g.muijzer@uva.nl

Editor: [Jana Jass]

#### Abstract

Haloalkaliphilic chemolithoautotrophic sulfur-oxidizing bacteria belonging to the genus Thioalkalivibrio are highly abundant in microbial communities found in soda lakes and dominant in full-scale bioreactors removing sulfide from industrial waste gases. Despite certain soda lakes being remote and unaffected by anthropogenic activities, haloalkaliphilic microorganisms, including Thioalkalivibrio strains, possess various antibiotic resistance genes. In this study, we investigated the impact of the antibiotic ampicillin on a co-culture of two Thioalkalivibrio species, Tv. thiocyanoxidans ARh2<sup>T</sup> and Tv. versutus AL2<sup>T</sup>, both experimentally and through in silico analysis of antibiotic resistance. Cell growth dynamics were monitored over time at increasing ampicillin concentrations using repand qPCR. Within ten days after the addition of ampicillin, the co-culture transitioned from a Tv. thiocyanoxidans ARh2<sup>T</sup>-dominated to a stable Tv. versutus AL2<sup>T</sup>-dominated culture. This shift was attributed to Tv. versutus AL2<sup>T</sup> displaying a lower susceptibility to ampicillin, making it more competitive. These results emphasize the potential implications of antibiotic pressure on microbial communities, where a resistant species can outcompete a stable co-culture. This study presents the first evidence of such dynamics in haloalkaliphilic chemolithoautotrophs. By understanding the antibiotic resistance and the competitive dynamics of haloalkaliphilic bacteria like Thioalkalivibrio, we can gain insights into their behaviour and stress response.

Keywords: antibiotics; haloalkaliphilic; population dynamics; soda lakes; sulfur-oxidizing bacteria; Thioalkalivibrio

# Introduction

Soda lakes are unique stable haloalkaline environments harbouring diverse microbial communities dominated by prokaryotes (Sorokin et al. 2014, Sorokin et al. 2015). These lakes can be found worldwide in areas with arid climates and with groundwater rich in sodium bicarbonate. High evaporation rates transform these lakes into highly sodium carbonate/bicarbonate-buffered, stable alkaline systems able to reach salt saturation (Jones et al. 1977, Grant et al. 1990, Jones et al. 1998). Despite these extreme conditions, the microbial communities are well adapted and flourish in these environments (Jones et al. 1998, Ochsenreiter et al. 2002, Mesbah et al. 2007, Lanzén et al. 2013), in which the chemolithoautotrophic sulfur-oxidizing bacteria of the genus Thioalkalivibrio are abundantly present (Sorokin et al. 2013, Vavourakis et al. 2016, Edwardson and Hollibaugh 2017, 2018, Zorz et al. 2019). In addition to soda lakes, Thioalkalivibrio strains are also important members of the microbial communities in the so-called "Thiopaq process", in which they sustainably remove sulfide from industrial waste gas through oxidation to elemental sulfur under haloalkaliphilic conditions (Sorokin et al. 2008, 2012). Recently, the antibiotic and metal "resistome" of the microbial communities of Lonar Lake, an anthropogenicpolluted alkaline lake in India, was described (Chakraborty et al. 2020a,b), for which Thioalkalivibrio has been significantly associated with antibiotic resistance genes (Chakraborty et al. 2020a).

The increased occurrence of antibiotic resistant bacteria and their associated genes in waterbodies due to anthropogenic pressure is an alarming phenomenon (Chakraborty et al. 2020b, Zhang et al. 2009, Vaz-Moreira et al. 2014, Yang et al. 2017, Su et al. 2020). However, antibiotic resistance as result of long-term evolution, existed long before anthropogenic pressure (D'Costa et al. 2011, Perry et al. 2016), protecting bacteria against their own antibiotic production and enabling them to outcompete other microbes in the ecosystem (Fajardo et al. 2009, Granato et al. 2019). Applying sub-inhibitory concentrations in humans, livestock or the environment poses a major threat to bacterial disease treatments with antibiotics, as it leads to an increase of antibiotic resistance amongst pathogens (Andersson and Hughes 2012). Alternatively, an antibiotic susceptible strain can be swept out of the population by an emerging resistant strain, which initially composed only a small fraction of the population, if an antibiotic concentration between the minimal inhibitory concentration (MIC) of the susceptible strain and the MIC of the resistant strain is applied. This concept is described as the "mutant selection window" hypothesis (Negri et al. 1994, 2000, Drlica and Zhao 2007).

Even though certain soda lakes are situated in remote regions without anthropogenic pressure, various haloalkaliphilic microor-

Received 5 February 2023; revised 26 October 2023; accepted 7 November 2023

© The Author(s) 2023. Published by Oxford University Press on behalf of FEMS. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (https://creativecommons.org/licenses/by/4.0/), which permits unrestricted reuse, distribution, and reproduction in any medium, provided the original work is properly cited.

ganisms from those environments possess a diverse spectrum of different antibiotic resistances (Pikuta et al. 2003, Boldareva et al. 2008, 2009a,b, Wu et al. 2010, Zargar et al. 2010, Kompantseva et al. 2012, Liu et al. 2012, Kiplimo et al. 2019). This has also been shown for strains of *Thioalkalivibrio* (Chakraborty et al. 2020a). With the interest to explore the antibiotic resistance capacity within *Thioalkalivibrio*, we studied here the dynamics of two *Thioalkalivibrio* species growing as a co-culture at increasing concentrations of ampicillin over a period of four weeks followed by four weeks without antibiotics. The dynamics in the co-culture were monitored with repetitive-element PCR (rep-PCR), which is a molecular fingerprint technique able to distinguish between the two *Thioalkalivibrio* species, and with quantitative PCR (qPCR) targeting strain-specific genes.

# Material and methods Cultivation and adaptation

Cultures of Tv. thiocyanoxidans ARh2<sup>T</sup> with low numbers of Tv. versutus AL2<sup>T</sup> were grown in 10 ml batch cultures at 100 rpm and 30°C with increasing concentrations of ampicillin for four weeks followed by four weeks without the addition of ampicillin. During the ampicillin adaptation phase, the culture was changed at each transfer to fresh medium with a doubled concentration in ampicillin, but it was also transferred again to fresh medium with the same ampicillin concentration. This was in case the culture would not be able to grow at the increased ampicillin concentration and needed more time to adapt at the lower concentration. The cultures started at 0.1 µg/ml ampicillin and were able to grow until 12.8  $\mu$ g/ml. No growth could be obtained with 25.6  $\mu$ g/ml. The experimental transfer schedule of the complete experiment can be found in Table S1 and is illustrated for the adaptation phase in Fig. S1. For each new culture, a 1:200 dilution of the previous culture was applied during the serial transfers. The medium was composed of 17.5 g/l Na<sub>2</sub>CO<sub>3</sub>, 13.9 g/l NaHCO<sub>3</sub>, 6.1 g/l NaCl, 1 g/l K<sub>2</sub>HPO<sub>4</sub>, 0.2 g/l MgCl<sub>2</sub>, 40 mM Na<sub>2</sub>S<sub>2</sub>O<sub>3</sub>, 5 mM NH<sub>4</sub>Cl and 1:1 000 trace metals (Pfennig and Lippert 1966), and the pH was adjusted to pH 9.8. Before inoculating the bacteria, a sterile solution of ampicillin (Roche Diagnostics GMBH, Mannheim, Germany) was added to the medium. Cultures with added ampicillin were prepared in triplicates and one culture without ampicillin was used as a reference. The MIC of both species was tested in triplicate using the same culture medium and the same ampicillin concentrations as for the adaptation experiment. Growth of the bacteria was followed daily by measuring the optical density in triplicates at 600 nm, and the average and the standard deviation of the three measurements were used in the data analysis.

#### **Repetitive-element PCR**

Samples were taken in the exponential phase at the end of each adaptation step and were pelleted by centrifugation at 15000 × g for 2 min. The supernatant was removed and the pellets were stored at  $-20^{\circ}$ C until further processing. Genomic DNA was extracted with the DNeasy Blood & Tissue Kit (Qiagen, Venlo, The Netherlands) following the supplier's instructions for Gramnegative bacteria.

Rep-PCR genomic fingerprinting was performed using the BOX A1R (5'-CTACGGCAAGGCGACGCTGACG-3') primer (Versalovic et al. 1994). The PCR protocol included an initial denaturation step at 94°C for 3 min, followed by 30 cycles of two denaturation steps of 94°C for 3 sec and 92°C for 3 sec, an annealing step of 46.4°C for 1 min, and an elongation step at 72°C for 8 min, and a final elonga-

tion step at 72°C for 8 min (Foti et al. 2006). The reaction included in a total reaction volume of 25  $\mu$ l, the Taq PCR Master Mix Kit 2x (Qiagen, Venlo, The Netherlands), 0.5  $\mu$ M primer, and 90 ng of template DNA. Rep-PCR products were visualized on a 1.5% (w/v) agarose gel stained with 0.6  $\mu$ g/ml ethidium bromide, which ran at 60 V for 8 h at 4°C. The GeneRuler DNA Ladder Mix (Thermo Fisher Scientific, Waltham, MA, USA) was used as a marker.

#### **Quantitative PCR**

The qPCR was performed in an ABI 7500 Real-Time PCR system (Applied Biosystems, Waltham, MA, USA) with strain-specific primers on the same template DNA as were used for the rep-PCR. The primers were designed with the Primer3 program (https: //primer3.ut.ee) and are listed with their target strain and genes in Table 1. The specificity of the primers was assessed in silico by BLAST and confirmed by a PCR on genomic DNA of Tv. versutus  $AL2^{T}$ , Tv. thiocyanoxidans  $ARh2^{T}$  and a 1:1 mixture of both species. The PCR protocol included an initial denaturation step at 94°C for 3 min, followed by 35 cycles of a denaturation at 94°C for 30 sec, an annealing at 60°C for 30 sec and an extension at 72°C for 15 sec, and a final extension at 72°C for 10 min in 25  $\mu$ l of reaction mix containing Taq PCR Master Mix Kit 2x (Qiagen, Venlo, The Netherlands), 0.5  $\mu\text{M}$  primer, and 10 ng DNA. PCR products were revealed by gel electrophoresis (Figure S2). The qPCR protocol included an initial denaturation at 95°C for 10 min, followed by 40 cycles of a denaturation at 95°C for 15 sec, an annealing at 60°C for 30 sec and an extension at 72°C for 30 sec. The reaction mix contained Maxima SYBR Green qPCR Master Mix (2x) (Thermo Fisher Scientific, Waltham, MA, USA), 0.3  $\mu$ M of forward and reverse primers, and 2 ng DNA in a total reaction volume of 25  $\mu$ l. ROX solution (Thermo Fisher Scientific, Waltham, MA, USA) was used to correct for well-to-well variation and melting curves were analysed for all measured samples to exclude non-specific PCR products. Each qPCR plate contained a reference sample (0 µg/ml-Ref) and a reference gene (16S rRNA) to overcome plate effects. For each individual sample, three technical replicates were measured, and the average and the standard error of the three measurements were used in the data analysis.

Individual amplification efficiencies (E) were calculated with LinRegPCR v.2018.0 (Ramakers et al. 2003, Ruijter et al. 2009) and were found between 1.8 and 2.0 with the exception at very low target numbers. The data was baseline corrected and threshold cycle ( $C_t$ ) values were calculated using LinRegPCR. Relative gene quantification was calculated using the initial culture before adaptation ( $T_0$ ) as reference sample and the 16S rRNA as reference gene according to the  $2^{-\Delta\Delta Ct}$  method (Livak and Schmittgen 2001). This type of quantification compares the changes in the target gene's copy number in a sample of interest to those in a reference sample (Livak and Schmittgen 2001). Negative control samples, in which the DNA template was replaced by water, did not show amplification.

#### In silico analysis on ampicillin resistance genes

The genomes of Tv. versutus  $AL2^{T}$  (NZ\_MVAR00000000) and Tv. thiocyanoxidans  $ARh2^{T}$  (NZ\_ARQK00000000) were previously sequenced and annotated. The gene sequences of both species were obtained from the NCBI RefSeq FTP server. Blastp analyses were performed to detect the presence of beta-lactamase genes in the genome of Tv. versutus  $AL2^{T}$  and Tv. thiocyanoxidans  $ARh2^{T}$ . For this, the beta-lactamase protein gene sequences of TEM, SHV, OXA and AmpC from other Gammaproteobacteria were used as references (Uniprot: Q6SJ61, Q0QXE5, Q5MD99, P0A9Z8, C7S9D0,

#### Table 1. Primers used in the qPCR.

Gene name*	Sequence (5´-3´) (Forward or Reverse)	Target strain (Locus tag)	Fragment length (bp)
16S rRNA	CGTGTGTGAAGAAGGCCTG (F) CCGGTGCTTCTTCTGTAGGT (R)	AL2 <sup>T</sup> (B0684_RS11480), ARh2 <sup>T</sup> (G372_RS0102255)	104
TcDH	AATACGTACTGCTGGCACCG (F) AGTTCCATCCAGCCAACCAC (R)	ARh2 <sup>T</sup> (G372_RS0106320)	96
PMOR	GTGCGTATCCTCTTCGACC (F) ATCACAGAACTGCCCGTGG (R)	AL2 <sup>T</sup> (B0684_RS04235)	102

\*TcDH encodes the thiocyanate: cytochrome c oxidoreductase; PMOR encodes the peptide-methionine (S)-S-oxide reductase.

P0A3M3, Q5EET9, D6NSM8). Moreover, the KEGG pathway map for beta-lactam resistance was analyzed for both strains and genes encoding for multidrug efflux pumps, RND family efflux transporters, AcrAB-TolC efflux pumps and MFS transporters were retrieved for analysis from the genomes of both species.

### **Results**

# Revealing ampicillin-driven species shifts in the co-culture

The rep-PCR patterns of a culture treated with increasing ampicillin concentrations and of the reference grown in the absence of ampicillin were followed in time (Fig. 1). The rep-PCR pattern of Tv. thiocyanoxidans ARh2<sup>T</sup> was present for the initial culture and for the culture containing 0.1  $\mu g/\text{ml}$  ampicillin. A mix of both species' rep-PCR fingerprint was seen for the culture at 0.2 µg/ml and at 0.4 µg/ml. The rep-PCR fingerprint then switched completely to the fingerprint of Tv. versutus AL2<sup>T</sup> at ampicillin concentrations of 0.8 µg/ml and higher. However, the reference culture incubated without ampicillin, did not experience a shift in pattern and maintained its Tv. thiocyanoxidans ARh2<sup>T</sup> pattern. Characteristic bands with the BOX A1R primer were found for Tv. thiocyanoxidans ARh2<sup>T</sup> at ~470 bp, ~520 bp, ~850 bp, ~950 bp, ~1100 bp,  $\sim$ 1850 bp and  $\sim$ 2200 bp, whereas for Tv. versutus AL2<sup>T</sup> they were found at ~320 bp, ~510 bp, ~650 bp, ~960 bp, ~1300 bp, ~1550 bp, and ~2300 bp.

The relative abundance of *Tv.* thiocyanoxidans  $ARh2^{T}$  and *Tv.* versutus  $AL2^{T}$  in the culture was monitored using qPCR amplification of strain-specific genes (Fig. 2). The gene encoding for thiocyanate: cytochrome *c* oxidoreductase (TcDH) was selected for *Tv.* thiocyanoxidans  $ARh2^{T}$  and the gene encoding the peptidemethionine (S)-S-oxide reductase (PMOR) for *Tv.* versutus  $AL2^{T}$  as they are single-copy genes and not present in the respective other species. The 16S rRNA gene is also only present as a single-copy gene in both species and was used as a reference for normalisation. The specificity of all three primers was evaluated as good after testing them in silico as well as by PCR using the genomic DNA of both species and a 1:1 mixture of DNA of both species (Figure S2).

The reference culture, which was incubated without ampicillin, maintained its initial composition of *Tv. thiocyanoxidans* ARh2<sup>T</sup> in the qPCR. However, cultures with increasing ampicillin concentration started to shift their composition from a *Tv. thiocyanoxidans* ARh2<sup>T</sup>-dominated culture as seen with 0 and 0.1 µg/ml ampicillin to a mixture of *Tv. thiocyanoxidans* ARh2<sup>T</sup> and *Tv. versutus* AL2<sup>T</sup> when the ampicillin concentration increased to 0.2 and 0.4 µg/ml between day three and eight. The original species *Tv. thiocyanoxidans* ARh2<sup>T</sup> was fully outcompeted by *Tv. versutus* AL2<sup>T</sup> after an increase to 0.8 µg/ml ampicillin sampled at day 10. The initial composition with *Tv. thiocyanoxidans* ARh2<sup>T</sup> did not recover after ten

successive incubations without ampicillin over a period of four weeks.

# Sensitivity of pure cultures to ampicillin

To understand the shift in species composition after incubation with the increasing concentrations of ampicillin, the sensitivity of Tv. versutus AL2<sup>T</sup> and Tv. thiocyanoxidans ARh2<sup>T</sup> was tested using the same concentrations as in the adaptation experiment (Figure S3). Tv. versutus  $AL2^{T}$  was able to grow with 0.1 µg/ml after two days and with 0.2 and 0.4  $\mu$ g/ml after three days lag period whereas Tv. thiocyanoxidans ARh2<sup>T</sup> showed only limited growth with ampicillin with 0.1 µg/ml after only three days. From a concentration of 0.8  $\mu$ g/ml and of 0.2  $\mu$ g/ml onwards, Tv. versutus AL2<sup>T</sup> and Tv. thiocyanoxidans ARh2<sup>T</sup>, respectively, were only able to grow after four days of incubation, which is likely due to degradation of ampicillin in the growth medium. Therefore, the respective MIC for ampicillin for Tv. versutus AL2<sup>T</sup> and Tv. thiocyanoxidans ARh2<sup>T</sup> are 0.8 µg/ml and 0.2 µg/ml. Chemical degradation of ampicillin in the culture medium was tested by inoculation of Thioalkalivibrio after one, two, three, or four days. Independently of when Thioalkalivibrio was added, visible growth only started after the fourth day when ampicillin was added to the medium (data not shown). In the absence of ampicillin, both species showed growth within one day. The OD<sub>600</sub> peak at the start of each Thioalkalivibrio culture resulted through the presence of elemental sulfur produced by the oxidation of thiosulfate. Its decrease after one day is linked to the depletion of sulfur due to complete oxidation to sulfate.

# Genomic exploration of ampicillin resistance genes in Thioalkalivibrio

Analysis of the genome sequences of Tv. versutus AL2<sup>T</sup> and Tv. thiocyanoxidans ARh2<sup>T</sup> was performed to retrieve their ampicillin and general antibiotic resistance capacity and to reveal differences. No beta-lactamase protein gene sequences of TEM, SHV, OXA and AmpC could be detected in either of the Thioalkalivibrio genomes using BLASTp against known protein sequences of other Gammaproteobacteria. The KEGG pathway map for beta-lactam resistance showed limited presence of genes in both species, but included penicillin-binding proteins and multidrug efflux pumps and Tv. versutus AL2<sup>T</sup> only possesses one supplementary gene (B0684\_RS10505), which could not be found in Tv. thiocyanoxidans ARh2<sup>T</sup> (Figure S4, Figure S5, Table S2). Both genomes were also screened for genes annotated as putative multidrug efflux pumps, putative RND family efflux transporters, putative AcrAB-TolC efflux pumps and putative MFS transporters (Table S2). Most genes could be retrieved in both genomes, but four genes are unique to Tv. thiocyanoxidans ARh2<sup>T</sup>, whereas eight to Tv. versutus AL2<sup>T</sup>. For Tv. thiocyanoxidans ARh2<sup>T</sup>, those genes are a multiple antibiotic resistance protein MarC (G372\_RS0106695), multidrug resistance efflux pump (G372\_RS0103685), multidrug efflux pump subunit AcrB (G372\_RS0103690) and an outer membrane



**Figure 1.** Rep-PCR patterns showing the dynamics of a co-culture of *Tv. thiocyanoxidans* ARh2<sup>T</sup> and *Tv. versutus* AL2<sup>T</sup> incubated with increasing ampicillin concentrations over time. The reference is a culture from the same starting culture as the adapted cultures, but without addition of ampicillin. NoAmp5 and NoAmp10 represent the adapted cultures after five and ten transfers without ampicillin, respectively. M represents the DNA marker.



**Figure 2.** Dynamics of the co-culture followed over time by qPCR with TcDH, which is specific to *Tv. thiocyanoxidans* ARh2<sup>T</sup> and PMOR, which is specific to *Tv. versutus* AL2<sup>T</sup>. Adapted culture 1, 2, and 3 represent triplicates from the same starting culture and were kept separate from each other throughout the experiment. The reference is a culture from the same starting culture as the adapted cultures, but without addition of ampicillin. The error bars depict the standard error of the average.

protein TolC (G372\_RS0109555). For *Tv. versutus*  $AL2^{T}$ , those include two multidrug efflux pump subunits AcrB (B0684\_RS03445, B0684\_RS12195), two RND family efflux transporter-MFP subunits (B0684\_RS03450, B0684\_RS09870), an outer membrane protein TolC (B0684\_RS04965), a multiple antibiotic resistance protein MarC (B0684\_RS07955), an outer membrane protein-multidrug efflux system (B0684\_RS08515), and a multidrug resistance efflux pump (B0684\_RS12200).

## **Discussion**

Our study supports the concept of the "mutant selection window" hypothesis. This concept states that a resistant subpopulation, which is initially only present as a minor fraction in the population, can emerge and outcompete a susceptible subpopulation from the population, if an antibiotic concentration between the MIC of the susceptible and the MIC of the resistant subpopulation is applied (Negri et al. 1994, 2000, Drlica and Zhao 2007). To prevent emergence of resistance from a pre-existing subpopulation, a "mutant prevention concentration" needs to be applied, which is a concentration higher than the MIC of the most resistant subpopulation (Drlica and Zhao 2007). This phenomenon is described based on two mutant subpopulations of a strain. However, the same principle can also be applied on two strains of a species, or on two species of a genus with the latter being the case in our study. The "mutant selection window" hypothesis was relativised by the study of Alexander and Maclean (2020). They predicted that at one-eighth of their MIC, individual resistant cells have only a less than 5% probability of detectable outgrowth. Moreover, they suggest that antibiotic concentrations below the MIC of the resistant strain might be efficient enough to eradicate the bacterial population when resistant mutants only make up a small fraction in the beginning (Alexander and Maclean 2020). However, it has been shown in our study as well as in multiple others that the selective antibiotic pressure can impact the population composition when a concentration below the MIC of the sensitive strain was applied. This provides a window of opportunity for resistant strains to outcompete the susceptible ones (Negri et al. 2000, Gullberg et al. 2011, Andersson and Hughes 2012, Hughes and Andersson 2012, Day et al. 2015).

Screening the composition of the co-culture with rep- and qPCR during the incubation with increasing concentrations of ampicillin revealed a switch in species dominance of the co-culture. Notably, the more resistant species Tv. versutus AL2<sup>T</sup>, which in the beginning of the experiment was only a minor component of the co-culture and not detectable by qPCR, outcompeted the more susceptible species Tv. thiocyanoxidans ARh2<sup>T</sup>. Within ten days and with an eight-fold increase of the ampicillin concentration, the resistant species became dominant in the co-culture and the sensitive species was no longer detectable. Growth profiles with different ampicillin concentrations explained this shift in dominance by a higher MIC for Tv. versutus AL2<sup>T</sup>. Competitivity of Tv. thio*cyanoxidans* ARh2<sup>T</sup> in the absence of ampicillin pressure as seen in the control culture could be explained on the other hand by the theory that the resistant strain has a higher fitness cost, for which reason the sensitive strain can outcompete the resistant strain (Gullberg et al. 2011).

The in silico analysis of both species' genomes did not detect any beta-lactamase genes and only revealed a limited number of genes associated with beta-lactam resistance. However, multiple putative multidrug efflux pump, putative RND family efflux transporters, putative AcrAB-TolC efflux pumps and putative MFS transporters, which have been previously associated to antibiotic resistance (Nikaido and Takatsuka 2009, Pérez et al. 2012, Nag and Mehra 2021), could be detected in both genomes with four genes specific to *Tv. thiocyanoxidans* ARh2<sup>T</sup> and eight to *Tv. versutus* AL2<sup>T</sup>. It is possible that amongst the specific genes in *Tv. versutus* AL2<sup>T</sup>, a gene could be present allowing its increase resistance to ampicillin compared to *Tv. thiocyanoxidans* ARh2<sup>T</sup>. However, their specific function and implication in ampicillin resistance still remains to be proven experimentally. Supplementary transcriptomic and proteomic analysis are needed to gain insights in the molecular mechanisms of ampicillin resistance in *Thioalkalivibrio* and to gain a better understanding of the increased ampicillin resistance of *Tv. versutus* AL2<sup>T</sup>.

Antibiotic treatments are associated with the evolution and the spread of resistances within microorganisms (Blair et al. 2014), as well as with shifts within the microbial community composition. These dynamics include changes in abundance and extinction of species in a given community (Coates et al. 2018, Cairns et al. 2020). Already a population composed by a single strain and subjected to bactericidal antibiotics showed population dynamics, but no stochastic dynamic was observed with bacteriostatic antibiotics (Coates et al. 2018). In a recent study on the impact of antibiotic perturbations on a complex microbial consortium, Cairns et al. (2020) found a repeatable community response. In their experiment, mock communities composed of 34 bacterial species were challenged with a pulsed antibiotic disturbance at different concentrations, with or without species immigration, to study the ecological community response. They proposed that this response could be linked to the antibiotic susceptibility and the growth rate of each given species. Next to species sorting, also antibiotic resistance evolution occurred in their experiment and its magnitude increased with increasing concentration of antibiotics. However, the genomic evolution seemed not to be relevant for the shortterm ecological dynamics in the studied community as the experiment showed high repeatability amongst the replicates (Cairns et al. 2020). In our study, a growing culture could not be obtained with ampicillin concentration higher than 12.5 µg/ml, which suggests a lack of rescuing genetic mutations in the final Tv. versutus AL2<sup>T</sup>-dominated co-culture.

Our results demonstrate how a stable population, which is dominated by a susceptible species can be exchanged by a stable population composed by a resistant species due to antibiotic pressure. To our knowledge it is the first time that the application of the "mutant application window" hypothesis is shown for haloalkaliphilic chemolithoautotrophs.

## Acknowledgment

We would like to thank Chiara de Jong, Mischa Tensen and Célia Roux for preliminary experiments on this subject. This research was financially supported by the ERC Advanced Grant PARASOL (No. 322551).

### Supplementary data

Supplementary data is available at FEMSMC Journal online.

Conflicts of interest: We do not have any conflict of interest.

# References

Alexander HK, Maclean RC. Stochastic bacterial population dynamics restrict the establishment of antibiotic resistance from single cells. Proc Natl Acad Sci USA 2020;**117**:19455–64. https://doi.org/10 .1073/pnas.1919672117.

- Andersson DI, Hughes D. Evolution of antibiotic resistance at nonlethal drug concentrations. *Drug Resist Updat* 2012;**15**:162–72. ht tps://doi.org/10.1016/j.drup.2012.03.005.
- Blair JMA, Richmond GE, Piddock LJV. Multidrug efflux pumps in gram-negative bacteria and their role in antibiotic resistance. Future Microbiology 2014;9:1165–77. https://doi.org/10.2217/fmb.14.6 6.
- Boldareva EN, Akimov VN, Boychenko VA et al. Rhodobaca barguzinensis sp. nov., a new alkaliphilic purple nonsulfur bacterium isolated from a soda lake of the Barguzin Valley (Buryat Republic, Eastern Siberia). Microbiology 2008;**77**:206–18. https://doi.org/10.1 134/S0026261708020148.
- Boldareva EN, Moskalenko AA, Makhneva ZK et al. Rubribacterium polymorphum gen. nov., sp. nov., a novel alkaliphilic nonsulfur purple bacterium from an Eastern Siberian soda lake. Microbiology 2009;**78**:732–40. doi:https://doi.org/10.1134/S0026261709060101.
- Boldareva EN, Tourova TP, Kolganova TV et al. Roseococcus suduntuyensis sp. nov., a new aerobic bacteriochlorophyll a-containing bacterium isolated from a low-mineralized soda lake of Eastern Siberia. Microbiology 2009;**78**:92–101. https://doi.org/10.1134/S002 6261709010123.
- Cairns J, Jokela R, Becks L *et al*. Repeatable ecological dynamics govern the response of experimental communities to antibiotic pulse perturbation. *Nat Ecol Evol* 2020;**4**:1385–94. https://doi.org/10.103 8/s41559-020-1272-9.
- Chakraborty J, Sapkale V, Rajput V et al. Shotgun metagenome guided exploration of anthropogenically driven resistomic hotspots within Lonar soda lake of India. Ecotoxicol Environ Saf 2020;194:110443. https://doi.org/10.1016/j.ecoenv.2020.110443.
- Chakraborty J, Sapkale V, Shah M et al. Metagenome sequencing to unveil microbial community composition and prevalence of antibiotic and metal resistance genes in hypersaline and hyperalkaline Lonar Lake, India. Ecol Indic 2020;110:105827. https://doi.or g/10.1016/j.ecolind.2019.105827.
- Coates J, Park BR, Le D et al. Antibiotic-induced population fluctuations and stochastic clearance of bacteria. eLife 2018;7:e32976. https://doi.org/10.7554/eLife.32976.
- D'Costa VM, King CE, Kalan L et al. Antibiotic resistance is ancient. Nature 2011;**477**:457–61. https://doi.org/10.1038/nature10388.
- Day T, Huijben S, Read AF. Is selection relevant in the evolutionary emergence of drug resistance? *Trends Microbiol* 2015;**23**:126–33. ht tps://doi.org/10.1016/j.tim.2015.01.005.
- Drlica K, Zhao X. Mutant selection window hypothesis updated. Clin Infect Dis 2007;44:681–8. https://doi.org/10.1086/511642.
- Edwardson CF, Hollibaugh JT. Metatranscriptomic analysis of prokaryotic communities active in sulfur and arsenic cycling in Mono Lake, California, USA. ISME J 2017;11:2195–208. https://doi.org/10.1038/ismej.2017.80.
- Fajardo A, Linares JF, Martinez JL. Towards an ecological approach to antibiotics and antibiotic resistance genes. Clin Microbiol Infect 2009;15:14–16. https://doi.org/10.1111/j.1469-0691.2008.02688.x.
- Foti M, Ma S, Sorokin DY *et al.* Genetic diversity and biogeography of haloalkaliphilic sulphur-oxidizing bacteria belonging to the genus *Thioalkalivibrio.* FEMS Microbiol Ecol 2006;**56**:95–101. https: //doi.org/10.1111/j.1574-6941.2006.00068.x.
- Granato ET, Meiller-Legrand TA, Foster KR. The evolution and ecology of bacterial warfare. *Curr Biol* 2019;**29**:R521–37. https://doi.org/10 .1016/j.cub.2019.04.024.
- Grant WD, Mwatha WE, Jones BE. Alkaliphiles: ecology, diversity and applications. FEMS Microbiol Lett 1990;**75**:255–69. https://doi.org/10.1016/0378-1097(90)90536-Y.

- Gullberg E, Cao S, Berg OG *et al*. Selection of resistant bacteria at very low antibiotic concentrations. PLoS Pathog 2011;**7**:e1002158. https://doi.org/10.1371/journal.ppat.1002158.
- Hughes D, Andersson DI. Selection of resistance at lethal and nonlethal antibiotic concentrations. *Curr Opin Microbiol* 2012;**15**:555– 60. https://doi.org/10.1016/j.mib.2012.07.005.
- Jones BE, Grant WD, Duckworth AW et al. Microbial diversity of soda lakes. Extremophiles 1998;2:191–200. https://doi.org/10.1007/s007 920050060.
- Jones BF, Eugster HP, Rettig SL. Hydrochemistry of the Lake Magadi basin, Kenya. *Geochim Cosmochim Acta* 1977;**41**:53–72. https://doi. org/10.1016/0016-7037(77)90186-7.
- Kiplimo D, Mugweru J, Kituyi S et al. Diversity of esterase and lipase producing haloalkaliphilic bacteria from Lake Magadi in Kenya. J Basic Microbiol 2019;59:1173–84. https://doi.org/10.1002/jobm.201 900353.
- Kompantseva EI, Komova AV, Novikov AA et al. Rhodovulum tesquicola sp. nov., a haloalkaliphilic purple non-sulfur bacterium from brackish steppe soda lakes. Int J Syst Evol Microbiol 2012;62:2962–6. https://doi.org/10.1099/ijs.0.035857-0.
- Lanzén A, Simachew A, Gessesse A *et al*. Surprising prokaryotic and eukaryotic diversity, community structure and biogeography of Ethiopian soda lakes. *PLoS One* 2013;**8**:e72577. https://doi.org/10 .1371/journal.pone.0072577.
- Liu YP, Wang YX, Li YX *et al*. Mongoliicoccus roseus gen. nov., sp. nov., an alkaliphilic bacterium isolated from a haloalkaline lake. Int J Syst Evol Microbiol 2012;**62**:2206–12. https://doi.org/10.1099/ijs.0. 035766-0.
- Livak KJ, Schmittgen TD. Analysis of relative gene expression data using real-time quantitative PCR and the 2<sup>-ΔΔCT</sup> method. *Methods* 2001;**25**:402–8. https://doi.org/10.1006/meth.2001.1262.
- Mesbah NM, Abou-El-Ela SH, Wiegel J. Novel and unexpected prokaryotic diversity in water and sediments of the alkaline, hypersaline lakes of the Wadi An Natrun, Egypt. Microb Ecol 2007;54:598-617. https://doi.org/10.1007/s00248-006-9193-y.
- Nag A, Mehra S. A major facilitator superfamily (MFS) efflux pump, SCO4121, from Streptomyces coelicolor with roles in multidrug resistance and oxidative stress tolerance and its regulation by a MarR regulator. Appl Environ Microb 2021;87:e02238–20. https://doi.org/ 10.1128/AEM.02238-20.
- Negri MC, Lipsitch M, Blásquez J et al. Concentration-dependent selection of small phenotypic differences in TEM β-lactamasemediated antibiotic resistance. Antibiotics 2000;2:237–55. https: //doi.org/10.3390/antibiotics2020237.
- Negri MC, Morosini MI, Loza E et al. In vitro selective antibiotic concentrations of β-lactams for penicillin-resistant Streptococcus pneumoniae populations. Antimicrob Agents Chemother 1994;**38**:122–5. https://doi.org/10.1128/AAC.38.1.122.
- Nikaido H, Takatsuka Y. Mechanisms of RND multidrug efflux pumps. Biochimica Et Biophysica Acta (BBA)—Proteins and Proteomics 2009;**1794**:769–81. https://doi.org/10.1016/j.bbapap.2 008.10.004.
- Ochsenreiter T, Pfeiffer F, Schleper C. Diversity of archaea in hypersaline environments characterized by molecular-phylogenetic and cultivation studies. Extremophiles 2002;**6**:267–74. https://doi. org/10.1007/s00792-001-0253-4.
- Pérez A, Poza M, Fernández A et al. Involvement of the AcrAB-TolC efflux pump in the resistance, fitness, and virulence of Enterobacter cloacae. Antimicrob Agents Chemother 2012;56:2084–90. https: //doi.org/10.1128/AAC.05509-11.
- Perry J, Waglechner N, Wright G. The prehistory of antibiotic resistance. Cold Spring Harb Perspect Med 2016;6:a025197. https://doi.or g/10.1101/cshperspect.a025197.

- Pfennig N, Lippert KD. Über das vitamin B<sub>12</sub>-bedürfnis phototropher Schwefelbakterien. Archiv Mikrobiol 1966;**55**:245–56. https://doi.or g/10.1007/BF00410246.
- Pikuta EV, Hoover RB, Bej AK et al. Tindallia californiensis sp. nov., a new anaerobic, haloalkaliphilic, spore-forming acetogen isolated from Mono Lake in California. Extremophiles 2003;7:327–34. https: //doi.org/10.1007/s00792-003-0326-7.
- Ramakers C, Ruijter JM, Lekanne Deprez RH et al. Assumptionfree analysis of quantitative real-time polymerase chain reaction (PCR) data. *Neurosci Lett* 2003;**339**:62–66. https://doi.org/10.1016/ s0304-3940(02)01423-4.
- Ruijter JM, Ramakers C, Hoogaars WMH et al. Amplification efficiency: linking baseline and bias in the analysis of quantitative PCR data. Nucleic Acids Res 2009;37:e45. https://doi.org/10.1093/ nar/gkp045.
- Sorokin DY, Banciu H, Robertson LA et al. (ed.), The Prokaryotes: Prokaryotic Physiology and Biochemistry. Berlin-Heidelberg: Springer-Verlag, 2013,530–55. https://doi.org/10.1007/978-3-642-30141-4.
- Sorokin DY, Banciu HL, Muyzer G. Functional microbiology of soda lakes. *Curr Opin Microbiol* 2015;**25**:88–96. https://doi.org/10.1016/j. mib.2015.05.004.
- Sorokin DY, Berben T, Melton ED *et al*. Microbial diversity and biogeochemical cycling in soda lakes. *Curr Opin Microbiol* 2015;**25**:88–96. https://doi.org/10.1016/j.mib.2015.05.004.
- Sorokin DY, Muntyan MS, Panteleeva AN et al. Thioalkalivibrio sulfidiphilus sp. nov., a haloalkaliphilic, sulfur-oxidizing gammaproteobacterium from alkaline habitats. Int J Syst Evol Microbiol 2012;62:1884–9. https://doi.org/10.1099/ijs.0.034504-0.
- Sorokin DY, Van Den Bosch PLF, Abbas B et al. Microbiological analysis of the population of extremely haloalkaliphilic sulfuroxidizing bacteria dominating in lab-scale sulfide-removing bioreactors. Appl Microbiol Biotechnol 2008;80:965–75. https://doi. org/10.1007/s00253-008-1598-8.
- Su S, Li C, Yang J et al. Distribution of antibiotic resistance genes in three different natural water bodies—A lake, river and sea. Int J

Environ Res Public Health 2020;**17**:552. https://doi.org/10.3390/ijer ph17020552.

- Vavourakis CD, Andrei AS, Mehrshad M *et al.* A metagenomics roadmap to the uncultured genome diversity in hypersaline soda lake sediments. *Microbiome* 2018;**6**:168. https://doi.org/10.1186/s4 0168-018-0548-7.
- Vavourakis CD, Ghai R, Rodriguez-Valera F et al. Metagenomic insights into the uncultured diversity and physiology of microbes in four hypersaline soda lake brines. Front Microbiol 2016;7:211. https://doi.org/10.3389/fmicb.2016.00211.
- Vaz-Moreira I, Nunes OC, Manaia CM. Bacterial diversity and antibiotic resistance in water habitats: searching the links with the human microbiome. FEMS Microbiol Rev 2014;38:761–78. https: //doi.org/10.1111/1574-6976.12062.
- Versalovic J, Schneider M, De Bruijn F et al. Genomic fingerprint of bacteria using repetitive sequence-based polymerase chain reaction. Methods Mol Cell Biol 1994;5:25–40.
- Wu XY, Zheng G, Zhang WW et al. Amphibacillus jilinensis sp. nov., a facultatively anaerobic, alkaliphilic bacillus from a soda lake. Int J Syst Evol Microbiol 2010;60:2540–3. https://doi.org/10.1099/ijs.0.01 8259-0.
- Yang Y, Xu C, Cao X *et al.* Antibiotic resistance genes in surface water of eutrophic urban lakes are related to heavy metals, antibiotics, lake morphology and anthropic impact. *Ecotoxicol* 2017;**26**:831–40. https://doi.org/10.1007/s10646017-1814-3.
- Zargar K, Hoeft S, Oremland R et al. Identification of a novel arsenite oxidase gene, arxA, in the haloalkaliphilic, arsenite-oxidizing bacterium Alkalilimnicola ehrlichii strain MLHE-1. J Bacteriol 2010;**192**:3755–62. https://doi.org/10.1128/JB.0 0244-10.
- Zhang XX, Zhang T, Fang HHP. Antibiotic resistance genes in water environment. Appl Microbiol Biotechnol 2009;82:397–414. https://do i.org/10.1007/s00253-008-1829-z.
- Zorz JK, Sharp C, Kleiner M et al. A shared core microbiome in soda lakes separated by large distances. Nat Commun 2019;10:4230. ht tps://doi.org/10.1038/s41467-019-12195-5.

Received 5 February 2023; revised 26 October 2023; accepted 7 November 2023

<sup>©</sup> The Author(s) 2023. Published by Oxford University Press on behalf of FEMS. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (https://creativecommons.org/licenses/by/4.0/), which permits unrestricted reuse, distribution, and reproduction in any medium, provided the original work is properly cited.