

**Machine learning based parameter sensitivity of regional climate models
a case study of the WRF model for heat extremes over Southeast Australia**

Reddy, P. Jyoteeshkumar; Chinta, Sandeep; Matear, Richard; Taylor, John; Baki, Harish; Thatcher, Marcus; Kala, Jatin; Sharples, Jason

DOI

[10.1088/1748-9326/ad0eb0](https://doi.org/10.1088/1748-9326/ad0eb0)

Publication date

2023

Document Version

Final published version

Published in

Environmental Research Letters

Citation (APA)

Reddy, P. J., Chinta, S., Matear, R., Taylor, J., Baki, H., Thatcher, M., Kala, J., & Sharples, J. (2023). Machine learning based parameter sensitivity of regional climate models: a case study of the WRF model for heat extremes over Southeast Australia. *Environmental Research Letters*, 19(1), Article 014010. <https://doi.org/10.1088/1748-9326/ad0eb0>

Important note

To cite this publication, please use the final published version (if applicable).
Please check the document version above.

Copyright

Other than for strictly personal use, it is not permitted to download, forward or distribute the text or part of it, without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license such as Creative Commons.

Takedown policy

Please contact us and provide details if you believe this document breaches copyrights.
We will remove access to the work immediately and investigate your claim.

LETTER • **OPEN ACCESS**


Machine learning based parameter sensitivity of regional climate models—a case study of the WRF model for heat extremes over Southeast Australia

To cite this article: P Jyoteeshkumar Reddy *et al* 2024 *Environ. Res. Lett.* **19** 014010

View the [article online](#) for updates and enhancements.

You may also like

- [Estimating reservoir evaporation using numerical weather prediction: Omo Gibe III reservoir in Ethiopia](#)
Abraham Loha Anebo, Tekalegn Ayele Woldesenbet and Gebiaw Teshome Ayele
- [Optimal Prediction of Atmospheric Turbulence by Means of the Weather Research and Forecasting Model](#)
Alohotsy Rafalimanana, Christophe Giordano, Aziz Ziad *et al.*
- [Performance comparison of solar radiation forecasting between WRF and LSTM in Gifu, Japan](#)
Jose Manuel Soares de Araujo



The Breath Biopsy® Guide
Fourth edition

FREE

DOWNLOAD THE FREE E-BOOK

BREATH BIOPSY

OWLSTONE MEDICAL

ENVIRONMENTAL RESEARCH
LETTERS

LETTER

OPEN ACCESS

RECEIVED
27 July 2023REVISED
14 November 2023ACCEPTED FOR PUBLICATION
21 November 2023PUBLISHED
30 November 2023

Original content from
this work may be used
under the terms of the
[Creative Commons
Attribution 4.0 licence](#).

Any further distribution
of this work must
maintain attribution to
the author(s) and the title
of the work, journal
citation and DOI.



Machine learning based parameter sensitivity of regional climate models—a case study of the WRF model for heat extremes over Southeast Australia

P Jyoteeshkumar Reddy^{1,*} , Sandeep Chinta² , Richard Matear¹ , John Taylor³ , Harish Baki⁴ ,
Marcus Thatcher⁵ , Jatin Kala^{6,7} and Jason Sharples^{7,8,9} ¹ Commonwealth Scientific and Industrial Research Organisation Environment, Hobart, TAS, Australia² Center for Global Change Science, Massachusetts Institute of Technology, Cambridge, MA, United States of America³ Commonwealth Scientific and Industrial Research Organisation Data61, Canberra, ACT, Australia⁴ Geosciences and Remote Sensing, Civil Engineering and Geosciences, TU Delft, Delft, The Netherlands⁵ Commonwealth Scientific and Industrial Research Organisation Environment, Melbourne, VIC, Australia⁶ Environmental and Conservation Sciences, Centre for Climate-Impacted Terrestrial Ecosystems, Harry Butler Institute, Murdoch University, Murdoch, WA, Australia⁷ Australian Research Council Centre of Excellence for Climate Extremes, University of New South Wales, Canberra, ACT, Australia⁸ School of Science, University of New South Wales Canberra, Canberra, ACT, Australia⁹ NSW Bushfire and Natural Hazards Research Centre, Sydney, NSW, Australia

* Author to whom any correspondence should be addressed.

E-mail: jyoteesh.papari@csiro.au**Keywords:** machine learning, sensitivity analysis, wrf, heat extremes, Southeast AustraliaSupplementary material for this article is available [online](#)**Abstract**

Heatwaves and bushfires cause substantial impacts on society and ecosystems across the globe. Accurate information of heat extremes is needed to support the development of actionable mitigation and adaptation strategies. Regional climate models are commonly used to better understand the dynamics of these events. These models have very large input parameter sets, and the parameters within the physics schemes substantially influence the model's performance. However, parameter sensitivity analysis (SA) of regional models for heat extremes is largely unexplored. Here, we focus on the southeast Australian region, one of the global hotspots of heat extremes. In southeast Australia Weather Research and Forecasting (WRF) model is the widely used regional model to simulate extreme weather events across the region. Hence in this study, we focus on the sensitivity of WRF model parameters to surface meteorological variables such as temperature, relative humidity, and wind speed during two extreme heat events over southeast Australia. Due to the presence of multiple parameters and their complex relationship with output variables, a machine learning (ML) surrogate-based global SA method is considered for the SA. The ML surrogate-based Sobol SA is used to identify the sensitivity of 24 adjustable parameters in seven different physics schemes of the WRF model. Results show that out of these 24, only three parameters, namely the scattering tuning parameter, multiplier of saturated soil water content, and profile shape exponent in the momentum diffusivity coefficient, are important for the considered meteorological variables. These SA results are consistent for the two different extreme heat events. Further, we investigated the physical significance of sensitive parameters. This study's results will help in further optimising WRF parameters to improve model simulation.

1. Introduction

Extreme weather events are occurring more frequently and intensely in recent years compared to the past across the globe (Masson-Delmotte *et al* 2021).

Among these extreme events, heatwaves and bushfires are challenging to society due to their substantial and persistent impact on human and natural systems (Perkins-Kirkpatrick and Lewis 2020, Abram *et al* 2021). Recent studies show that heatwaves

are intensifying globally and regionally in Australia, underscoring the urgent need for efficient mitigation and adaptation techniques (Perkins-Kirkpatrick and Lewis 2020, Reddy *et al* 2021). Due to its frequent exposure to heatwaves, bushfires, and other extreme events, Southeast Australia, in particular, has been identified as a region with a greater susceptibility to the effects of climate change (Bureau of Meteorology and CSIRO 2020, Sharples *et al* 2016, Di Virgilio *et al* 2019a). The increasing frequency and intensity of bushfires in the region have been linked to climate change, further aggravating the risks and consequences of these events (Sharples *et al* 2016, Canadell *et al* 2021). Considering these events have disastrous impacts, accurately simulating heat extremes using numerical weather prediction (NWP) models is critical to support appropriate planning of adaptation and mitigation efforts.

The accuracy of a simulation from an NWP model depends on the exactness of the initial conditions and the ability of the model to represent the physical processes that occur in the atmosphere (Bjerknes 1910). These physical processes occur at different spatial and temporal scales. Due to limited computational resources, the model cannot accurately represent the physical processes at all scales. Certain physical processes occur at smaller scales than those captured by the model grid. Moreover, even at scales equivalent to the grid resolution, some processes have complex mechanisms that are computationally intensive or intricate to be represented explicitly. Parameterizations address this shortcoming by offering simplified, computationally efficient representations of complex physical processes, such as convection, cloud formation, soil moisture transport, and boundary layer physics, to name a few. Parameterization schemes employ mathematical equations to capture these complex physical processes and relate these partially represented processes to the large-scale processes that the model resolves. These equations are typically based on empirical relationships or theoretical understanding derived from observations or detailed process studies (Kain 2004, Hong and Lim 2006). Numerous tuneable parameters are frequently included in the parameterization schemes to simulate atmospheric processes. These parameters are typically constants and exponents in the model equations, and their default values are specified by the scheme developers based on experimental or theoretical investigations (Hong *et al* 2004). Therefore, accurate specification of these parameters within the model's physical representation is crucial for the reliability of model simulations (Yang *et al* 2012, Di *et al* 2015).

The Weather Research and Forecasting (WRF) model is an NWP model widely used as a weather forecasting tool and for dynamical downscaling purposes across the globe. Researchers and

meteorological organizations have adopted the model extensively due to its adaptability, flexibility, and high-resolution capabilities (Skamarock *et al* 2021, Evans *et al* 2014). The WRF model has been used in numerous studies in simulating extreme events like heatwaves and extreme bushfires in Southeast Australia (Kala *et al* 2015, 2023, Di Virgilio *et al* 2019a). The WRF model has multiple options for parameterization schemes for each physical process. Different combinations of parameterization schemes give the user a choice of model configurations (Skamarock *et al* 2021). Previous studies have examined the sensitivity of various physics parameterization schemes in simulating extreme events over Southeast Australia (Evans *et al* 2012, 2014, Ji *et al* 2022) but no studies have explored the sensitivity of parameters within the physics schemes. As discussed earlier, the values of parameters in these physics parameterization schemes play a crucial role in model fidelity. Therefore, calibration of these parameter values using observations could help improve the model simulation (Duan *et al* 2017, Chinta and Balaji 2020, Baki *et al* 2022a).

Parameter estimation can be a complex process due to several challenges. With many tuneable parameters (often in tens to hundreds), the number of model simulations required to calibrate these parameters increase exponentially as the number of tuneable parameters increases. Furthermore, multiple meteorological variables, such as humidity, wind speed, and temperature, to name a few, must be considered simultaneously during calibration. The parameter calibration process requires enormous computational resources. This challenge can be overcome by first implementing sensitivity analysis (SA) to identify the parameters significantly influencing the model output variables of interest. This approach helps reduce the number of parameters that need calibration, ultimately improving the accuracy of the variables of interest in the model output (Quan *et al* 2016, Chinta *et al* 2021). SA investigates how variations in the model output can be attributed to different sources of uncertainty in the model's input parameters (Saltelli 2002). Several studies have implemented SA to WRF model parameters to identify the parameters that influence the output variables (Yang *et al* 2012, Di *et al* 2015, 2017, 2018, Ji *et al* 2018, Wang *et al* 2020, Baki *et al* 2022b).

This study aims to identify the WRF model parameters that most affect the different model output meteorological variables related to heat extremes (i.e. heatwaves and fire weather). We do this by focusing on Southeast Australia during two extreme fire weather and heatwave events. This study is organized as follows; section 2 introduces the surrogate model based global SA technique and describes the data, events selected, and WRF model parameters for the SA. Section 3 presents the SA results and a detailed

discussion of the physical significance of identified sensitive parameters. Section 4 summarises the conclusions from this study.

2. Methods

2.1. Machine learning (ML) surrogate-based

Sobol SA

SA is broadly classified into local and global categories. The local SA is performed by exploring how the perturbed input parameters around specific reference values influence the output(s). However, global SA explores the entire feasible parameter space and highlights each parameter's total effects, including their interactive effects on the model output. The local SA method is widely used because of its limited computational requirements (Rakovec *et al* 2014). However, if the input parameters interact and have a non-linear influence on the output(s), the local SA method will be substantially biased and underestimate the importance of parameters (Saltelli *et al* 2019). The considered WRF model parameters are expected to have complex interactions and non-linearly influence the outputs (Di *et al* 2015, Quan *et al* 2016, Chinta *et al* 2021, Baki *et al* 2022b). Hence, in this study, we use the global SA method. There are many global SA techniques, including Morris One-at-A-Time, Multivariate adaptive regression splines, Fourier amplitude sensitivity test, and Sobol SA. Out of these global SA methods, previous WRF parameter SA studies recommend the Sobol SA method because of its accuracy, capacity to address interaction effects, discontinuities, and possible non-linear effects of the parameters on the output variables (Wang *et al* 2020, Baki *et al* 2022b).

Sobol SA is based on Analysis of Variance decomposition, also known as Hoeffding-Sobol decomposition, which states that a functional output variance can be expressed as the combination of variances of the output function contributed by individual parameters, interactions of parameter pairs, and so on in the increasing dimensionality (Sobol' 2001, Saltelli *et al* 2010). Based on the definition of variance as a measure of likeliness, the Sobol sensitivity indices are estimated as the ratios of variances contributed by the parameter interactions with respect to the total variance. The computation of sensitivity indices requires functional evaluations over a large number of input samples, which is computationally very expensive for the WRF model. However, it is feasible to generate the thousands of output samples required for the SA by employing a surrogate ML model that has been well trained on a limited set of WRF model runs. Among the various ML models available for use as surrogate models, Gaussian process regression (GPR) has been identified as one of the best suitable choices in such scenarios (Ji *et al* 2018, Baki *et al* 2022b).

GPR is a probabilistic and non-parametric ML technique that is accurate enough to predict and model unknown functions. The GPR is built upon the Gaussian Process, which is 'a collection of random variables, any finite number of which have consistent Gaussian distributions' (Williams and Rasmussen 1995, 2006). For any vector of random input variables $x = \{x_1, x_2, \dots, x_n\}$, the corresponding latent functions $f(x) = \{f(x_1), f(x_2), \dots, f(x_n)\}$ form a joint multivariate Gaussian distribution governed by a mean function and kernel function. Based on this, the GPR can be explained in four stages. First, assuming a mean and a kernel function, a collection of random output latent functions is obtained over random input variables, known as a Gaussian prior distribution. Second, provided a set of observed training samples and assuming the targets follow a Gaussian distribution, the hyperparameters of the kernel function are obtained by maximizing the log-likelihood of the target's Gaussian distribution. Third, with the optimized hyperparameters, a posterior distribution can be modelled by conditioning the observations for a new set of test points. Finally, the posterior distribution can be used to make predictions of any set of new test points and keep the model updated.

2.2. WRF model configuration


This study employs the Advanced Research WRF (WRF-ARW) model version 4.4 for the numerical simulations (Skamarock *et al* 2021). The model domain encompassing southeast Australia consists of 206×181 grid points with a horizontal resolution of 12 km (figure S1). The model has 40 σ vertical levels with the top layer set at the 50 hPa atmospheric level. The time integration is performed with a time step of 72 s. The European Centre for Medium-Range Weather Forecast Reanalysis 5th generation data set (ERA5) (Hersbach *et al* 2020) at 0.25° horizontal resolution with a six-hourly interval are used as the initial and lateral boundary conditions. The WRF model output data is stored at hourly intervals. The parameters and physics schemes are adapted based on previous studies (Di *et al* 2015, Quan *et al* 2016, Chinta *et al* 2021, Baki *et al* 2022b). The similar WRF physics parametrisation scheme combination performed reasonably well in simulating the climatology and extreme temperatures over the study domain (Evans and McCabe 2010, Kala *et al* 2015). A total of 24 adjustable model parameters are identified across seven different physics schemes. These 24 parameters are chosen because they have documented ranges. The details are presented in table 1.

2.3. Experimental setup

In this study, we select the two heat extremes where heatwaves and extreme fire weather days are observed

Table 1. The WRF model physics schemes used in this study with corresponding adjustable parameters in the schemes and their respective default values and ranges. These default values and ranges are adapted from previous studies (Di *et al* 2015, Quan *et al* 2016, Chinta *et al* 2021, Baki *et al* 2022b).

Physics scheme	Parameter	Default	Range	Description
Surface layer-MM5 Monin–Obukhov Scheme (Jiménez <i>et al</i> 2012)	P1	2.4×10^{-5}	1.2×10^{-5} – 5×10^{-5}	The parameter for heat/moisture exchange coefficient (s m^{-2})
	P2	0.0185	0.01–0.037	The coefficient for converting wind speed to roughness length over water
	P3	1	0.5–2	Scaling related to surface roughness
	P4	0.4	0.35–0.42	Von Kármán constant
Cumulus-Kain-Fritsch Eta scheme (Kain 2004)	P5	1	0.5–2	The multiplier for downdraft mass flux rate
	P6	1	0.5–2	The multiplier for entrainment mass flux rate
	P7	150	50–350	Starting height of downdraft above USL (hPa)
	P8	2700	1800–3600	Average consumption time of CAPE (s)
	P9	5	3–12	The maximum turbulent kinetic energy value in sub-cloud layer ($\text{m}^2 \text{s}^{-2}$)
Microphysics-WSM 6-class scheme (Hong and Lim 2006)	P10	14900	8000–30000	Scaling factor applied to ice fall velocity (1 s^{-1})
	P11	8×10^6	5×10^6 – 12×10^6	Intercept parameter of rain (1 m^{-4})
	P12	5×10^{-4}	3×10^{-4} – 8×10^{-4}	The limited maximum value for the cloud-ice diameter (m)
	P13	0.55	0.35–0.85	Collection efficiency from cloud to rain auto conversion
Shortwave radiation-Dudhia scheme (Dudhia 1989)	P14	1×10^{-5}	0.5×10^{-5} – 2×10^{-5}	Scattering tuning parameter ($\text{m}^2 \text{ kg}^{-1}$)
Longwave radiation-RRTM scheme (Mlawer <i>et al</i> 1997)	P15	1.66	1.55–1.75	Diffusivity angle for cloud optical depth computation
Land surface-unified Noah land surface model scheme (Chen and Dudhia 2001, Tewari <i>et al</i> 2004)	P16	1	0.5–2	The multiplier for hydraulic conductivity at saturation
	P17	1	0.5–2	The multiplier for the saturated soil water content
	P18	1	0.5–2	The multiplier for minimum soil suction
	P19	1	0.5–2	The multiplier for Clapp and hornberegger ‘b’ parameter
Planetary boundary layer-Yonsei University (YSU) scheme (Hong <i>et al</i> 2006)	P20	0.3	0.15–0.6	Critical Richardson number for boundary layer of water
	P21	0.25	0.125–0.5	Critical Richardson number for boundary layer of land
	P22	2	1–3	Profile shape exponent for calculating the momentum diffusivity coefficient
	P23	6.8	3.4–13.6	Coefficient for Prandtl number at the top of the surface layer
	P24	15.9	12–20	Counter gradient proportional coefficient of non-local flux of momentum



over southeast Australia for the numerical simulations. One is the southeast Australian heatwave event starting in the last week of January and extending to the second week of February 2009 (Engel *et al* 2013). The 2009 event is simulated for 13 d covering the period of extreme heat, i.e. from 26th January 12 UTC to 08th February 12 UTC. Another event is the heat extremes during the mid to end of December 2019/20 period. The 2019 event is simulated for 15 d covering heatwave and extreme fire weather days, i.e. from 16th December 12 UTC to 31st December 12 UTC. A 36 h model spin-up is considered for both events. The simulation results are evaluated against the Bureau of Meteorology Atmospheric high-resolution Regional Reanalysis for Australia (BARRA2) data (Su *et al* 2022). BARRA2 provides hourly data at 12 km horizontal resolution over the Australian region.

This study conducts parameter SA for the critical meteorological variables of heat extremes such as temperature, relative humidity, and wind speed during both events (2009 and 2019). Firstly, a Quasi-Monte Carlo Sobol sequence design is employed to generate 256 parameter samples using the Uncertainty Quantification Python Laboratory package as recommended by the previous studies

(Wang *et al* 2020, Baki *et al* 2022b). Next, we perform the WRF simulations with the 256 generated parameter samples for each event. A sample size of 256 is selected based on the sample size sensitivity experiments (figure S2). Then we calculate each meteorological variable's root mean square error (RMSE) between WRF simulations and BARRA2 data. The RMSE of 256 WRF runs for both the events is presented in figure 1. The RMSE is normalised with min-max normalisation for each selected event, respectively. The 256 WRF runs with perturbed parameters are broadly consistent for both the events with minor differences which are expected due to slightly different event characteristics (figure 1).

We then train the GPR model with parameter samples as input and RMSE as output for each of the considered meteorological variables. The GPR model is evaluated against the WRF RMSE using a K -fold cross-validation technique. Here the entire dataset is divided into K -folds ($K = 8$). The GPR model is trained with $(K-1)$ folds; the left-out K th fold is used as test data. It is iterated over all the folds, and each test set's GPR predictions are stacked. The accuracy of the GPR model is measured with the goodness of fit (R^2) metric by comparing the GPR predictions with WRF

data. This shows that GPR has good accuracy with high R^2 values for each considered meteorological variable (figure S3). Consequently, the trained GPR model is used for predicting the RMSE of the 51 200 parameter samples. These parameter samples are generated using the Sobol sequence design, following previous studies (Wang *et al* 2020, Baki *et al* 2022b). Finally, Sobol sensitivity indices are calculated based on these 51 200 samples, thereby providing insights into the relative importance of the parameters. The detailed flowchart of ML surrogate based Sobol SA methodology is presented in supplementary figure S4. Further, to improve the robustness of the ML surrogate-based Sobol' SA results, following the previous studies (Wang *et al* 2020, Baki *et al* 2022a), we have also performed the response-surface-based SA with multivariate adaptive regression splines (MARS) using the 256 WRF sample. More details are presented in supplementary material.

3. Results and discussion

3.1. Parameter SA of heat extremes

The parameter SA is performed for the hourly surface (2 m) air temperature (T), surface (2 m) relative humidity (Rh), and surface (10 m) wind speed (Ws) during the two case studies. Both the first- and higher-order Sobol sensitivity indices of parameters for the corresponding meteorological variables are presented in figure 2. The first-order sensitivity indices represent the effect of variations in a single parameter, and higher-order sensitivity indices represent the interaction effects of changes in multiple parameters on the output (Sobol' 2001, Saltelli 2002, Saltelli *et al* 2010). The SA results are presented individually for the considered extreme heat events (2009 and 2019) and for the combined data in figure 2. It is observed that the SA results are mostly consistent across both events and also for the combined data. However, minor variations in the degree of sensitivity of the important parameters between the events were observed (figure 2). These variations in the magnitude of sensitivity could be due to the slightly different event characteristics (figure 1).


SA results suggest that parameters P14 (scattering tuning parameter), P17 (multiplier of saturated soil water content), and P22 (profile shape exponent in the momentum diffusivity coefficient) most affect the considered meteorological variables (T , Rh , and Ws). For T , only two parameters, P14 and P17, are most important, and their sensitivity is mostly first-order. Their higher-order sensitivity is minor compared to the first-order effects on T (figure 2(a)). This means that P14 and P17 individually affect the T to a large extent, and the higher-order effects are mostly due to the interactions between them (not shown). Similar to T , P14 and P17 are the most influential parameters for Rh , and in addition to these, P22 has some minor

influence on Rh . Again, for Rh , the sensitive parameters (P14, P17, and P22) influence is mostly first-order (figure 2(b)). In contrast to T and Rh , for Ws , P22 is equally or slightly more important as the P14 and P17 parameters (figure 2(c)). Results show that the higher-order sensitivity of P14 and P17 for Ws is considered to be as important as first-order sensitivity. This higher-order sensitivity of P14 and P17 for Ws is partly due to their second-order interactions (not shown). Several other parameters, P6, P10, P12, P21, P23, and P24, show very minor effects on Ws (figure 2(c)).

Similar to Sobol' SA, the MARS SA is performed on the 2009, 2019 events and combined data but only using the 256 WRF sample. MARS SA results are consistent with the Sobol' SA (figures 2 vs S5). Further, we validate the ML surrogate-based Sobol SA results by examining one additional extreme heat event i.e. 2013 January heatwave (for more details on the event refer to supplementary material). The 2013 and the combined data of all three events SA results are consistent with the 2009 and 2019 events SA results (figures 2 vs S6). These results further increase the confidence of SA results of the WRF model for heat extremes over southeast Australia.

3.2. The physical significance of sensitive parameters

Parameter P14 is the scattering tuning parameter in the Dudhia shortwave radiation scheme, which is related to the scattering under clear sky conditions. This parameter directly influences the incoming shortwave radiation reaching the Earth's surface by affecting the scattering attenuation (Dudhia 1989, Montornès Torrecillas *et al* 2015). The solar radiation is inputted to the land surface parameterisation scheme, which then transforms to surface energy fluxes (sensible and latent heat). The surface fluxes regulate the mixing in the planetary boundary layer through thermals, affecting the horizontal wind speed (Oke 2002). Therefore, T , Rh , and Ws are sensitive to P14. When P14 is low, scattering is attenuated, leading to an increase in incoming shortwave radiation, which then heats the surface and increases the surface temperature. Our results are consistent with this: as P14 decreases, the mean T values increase (figure 3(a)). This increase in mean T is mainly due to the increase in daily maximum temperature (T_{\max}) (figure 3(b)). P14's influence on daily minimum temperature (T_{\min}) is very minimal because the parameter is related to the scattering of shortwave radiation, which is absent during night time (figure 3(c)). Similar to T , P14 influences Rh (figures 3(d)–(f)). As P14 decreases, the mean Rh and mean daily minimum relative humidity (Rh_{\min}) decrease; however, the mean daily maximum relative humidity (Rh_{\max}) is mostly unaffected (figures 3(d)–(f)). For Ws , lower P14 results in stronger winds (both the mean and



daily maximum wind speed ($W_{s_{\max}}$) and vice versa (figures 3(g), (i), (j), and (l)).

Parameter P17, the multiplier of saturated soil water content or soil porosity in the Unified Noah land surface scheme, is the other most influential parameter for T , Rh , and Ws . P17 regulates the transmission rate of moisture and heat transport in the soil (Chen and Dudhia 2001, Tewari *et al* 2004). This

implicitly affects the rate of heat transfer and water vapour exchange between the land surface and the atmosphere (Tewari *et al* 2004, Chinta *et al* 2021). Smaller values of P17 implies less porosity, which means more dense soil with high soil thermal inertia. The high thermal inertia means lower day time temperatures and higher night time temperatures (Quan *et al* 2016, Fonseca *et al* 2019, Temimi *et al* 2020).




Figure 3. The mean of surface meteorological variable ((a) hourly temperature (T ; $^{\circ}\text{C}$), (b) daily maximum temperature (T_{max} ; $^{\circ}\text{C}$), (c) daily minimum temperature (T_{min} ; $^{\circ}\text{C}$), (d) hourly relative humidity (Rh), (e) daily minimum relative humidity (Rh_{min}), (f) daily maximum relative humidity (Rh_{max}), (g)–(i) hourly wind speed (Ws ; m s^{-1}), and (j)–(l) daily maximum wind speed (Ws_{max} ; m s^{-1})) of 256 WRF runs with a different set of parameter values for the combined data of both events. The x - and y -axis represent the respective sensitive parameters of corresponding meteorological variables (parameters P17 and P14 for temperature (a)–(c), relative humidity (d)–(f), and wind speed (g) and (j); P17 and P22 for wind speed (h) and (k); P22 and P14 for wind speed (i) and (l)). The color represents the mean values of the respective meteorological variables.

This is due to an increase in conduction which means faster heat transfer between the surface and subsurface, leading to a slower rate of surface warming (cooling) during the day (night). The slow rate of day time (night time) surface warming (cooling) results in low (high) daily maximum (minimum) temperatures. Our results are consistent with this and show that at low P17, mean T_{max} is lower and mean T_{min} is higher, compared to those during high P17 (figures 3(b) and (c)). Similar to T_{max} and T_{min} , as P17 decreases, the mean Rh_{min} increases; however, the mean Rh_{max} decreases (figures 3(e) and (f)). Overall, low P14 and low P17 results in high mean T and low mean Rh (figures 3(a) and (d)).

The parameter P22 is the profile shape exponent associated with the calculation of the momentum

diffusivity coefficient in the boundary layer scheme (Hong et al 2006). P22 governs the turbulent mixing in the boundary layer affecting the height of the maximum diffusivity and leading to the influence of the horizontal wind speed (Oke 2002, Di et al 2015, Baki et al 2022b). As P22 increases, the height of momentum diffusivity declines and results in increasing (decreasing) momentum diffusivity coefficient below (above) the maximum height. This means high P22 can lead to weak winds in low levels and strong winds in the upper levels of the atmosphere, respectively (Yang et al 2019). Our results show that lower P22 values are associated with stronger mean wind speeds (both Ws and Ws_{max}) and vice versa (figures 3(h), (i), (k), and (l)).




Figure 4. Spatial plot of average daily maximum temperature (T_{max} ; °C) (a), daily minimum relative humidity (Rh_{min}) (e), and daily maximum wind speed (Ws_{max} ; m s⁻¹) (i) during all days of both selected events (2009 and 2019) using the BARRA2 data. Comparison of the WRF default parameters run (default), best (respective minimum RMSE of corresponding hourly meteorological variable data out of 256 runs) parameters run (best (256)), and worst (respective maximum RMSE of corresponding hourly meteorological variable data out of 256 runs) parameters run with respect to BARRA2 data for the considered meteorological variables. The mean bias of T_{max} (b)–(d), Rh_{min} (f)–(h), and Ws_{max} (j)–(l) between default, best (256), and worst (256) runs with respect to BARRA2.

3.3. Comparing the default, best, and worst parameter combination results

The SA results show that only three parameters most affect the considered meteorological variables. These three sensitive parameters can be further optimised to improve WRF model simulation of critical meteorological variables for heat extremes such as T_{max} , Rh_{min} , and Ws_{max} . The calibration of sensitive parameters with a multi-objective optimisation technique is a complicated task, which is outside the present study's scope but can be done in a future study. However, the 256 WRF simulations for the SA can provide us with an understanding of how different parameter sets, compared to the default values, can improve (or amplify) the model biases of the selected output variables. For this purpose, we compare the results of default, best, and worst parameter combinations out of 256 WRF runs, as illustrated in figure 4. The parameter set with minimum (maximum) RMSE of corresponding hourly meteorological variable data (T , Rh , and Ws) out of 256 WRF runs is chosen as best (worst).

Figure 4 presents the comparison of average T_{max} , Rh_{min} , and Ws_{max} during all days of both events (2009 and 2019) for the default, best, and worst parameter sets with respect to BARRA2 data. The default simulation results suggest that T_{max} is under-predicted by around 2 °C–4 °C and over-predicted both the Rh_{min} (by about 0.06) and Ws_{max} (by approximately 2–4 m s⁻¹) compared to the BARRA2 data over much of the study domain. These changes are more

substantial over the eastern coastal strip (figures 4(b), (f), and (j)). These results are broadly consistent with the previous studies, which showed the cold biases in the default parameter WRF model simulated T_{max} compared to observations over the study region (Kala *et al* 2015, Di Virgilio *et al* 2019b, Ji *et al* 2022). However, the best parameter combination results suggest that the simulation of T_{max} , Rh_{min} , and Ws_{max} is improved over much of the study domain compared to the default run (figures 4(c), (g), and (k)). The improvements can be clearly seen over the regions of T_{max} greater than 40 °C (compare figures 4(b) and (c)). Conversely, T_{max} , Rh_{min} , and Ws_{max} biases are amplified for the worst parameter set compared to the default simulation (figures 4(d), (h), and (l)). These results are consistent even when only the extremely hot day of each event is considered separately (figures S7 and S8). Overall, these results suggest that tuning the most sensitive parameters can further improve model simulation of meteorological variables of interest.

3.4. Limitations

We would like to emphasize that this study has a few caveats and limitations that should be noted. The parameter SA results are mostly dependent on regional conditions and could be on the type of the selected event (here heat extremes) (Di *et al* 2015, Quan *et al* 2016, Ji *et al* 2018, Chinta *et al* 2021). Hence, the SA results of this study may not be precisely applicable to other regions or other kinds of

extreme events (for example, wet extremes). Testing the validity of this study's SA results for different extreme event types is beyond the scope of the present study.

Another limitation is that we have conducted the SA for only 24 parameters which may not be the complete list of all tuneable parameters in the WRF model. Nevertheless, this study is the first to systematically assess the sensitivity of WRF model parameters to considered meteorological variables during extreme heat events over southeast Australia. This study's results are mostly relevant for short simulations of these extremes, not necessarily for long-term regional climate model simulations. Nonetheless, our results will provide significant guidance for future research on parameter calibration of sensitive parameters for improving the WRF model simulation of heat extremes over southeast Australia. Future studies can use the present study's methodology or a similar one to conduct parameter SA of different regional atmosphere models in various regions for a variety of extreme events.

4. Conclusions

Studies using regional atmospheric modelling systems to better understand extreme heat events such as heatwaves and bushfires rarely examine the model's sensitivity to tuneable parameters within parameterization schemes. This is the first study to address this issue over southeast Australia, a region that is highly susceptible to such extremes. We have investigated the sensitivity of 24 WRF tuneable parameters from seven different physics parameterisation schemes to the meteorological variables critical to the heat extremes over southeast Australia. The ML surrogate-based Sobol SA method is used to conduct the parameter SA during the two extreme heat events (2009 and 2019) in southeast Australia. Results show that only three parameters, P14 (scattering tuning parameter), P17 (multiplier of saturated soil water content), and P22 (profile shape exponent in the momentum diffusivity coefficient), are most important for the considered meteorological variables (T, Rh, and Ws) during heat extremes in southeast Australia. These results are consistent for both the considered events. These parameters are in the shortwave radiation, land surface, and planetary boundary layer schemes.

Further, we examined the physical significance of sensitive parameters on how they affect the simulation of considered meteorological variables. In conclusion, results showed that both low values of P14 and P17 simulate high T and low Rh, whereas low P14 and low P22 produce strong winds. Overall, this study's results suggest that only three out of 24 parameters needed to be considered for improving the simulation of heat extremes over southeast Australia. Future studies can focus on optimising these three

sensitive parameters for improving the simulation of heat extremes over southeast Australia using multi-objective optimisation techniques. Our overall methodology will also be helpful for studies focussing on extreme heat events elsewhere.

Data availability statement

The data that support the findings of this study are openly available at the following URL/DOI: <https://doi.org/10.24381/cds.adbb2d47>.

Acknowledgments


The authors would like to thank Chun-Hsu Su for providing the BARRA2 data. We acknowledge James Risbey's suggestion for improving the figure 4. We would like to thank National Computing Infrastructure (NCI) Australia for providing computational resources. The authors would like to acknowledge the funding support of the CSIRO and Australian Climate Service.

ORCID iDs

P Jyoteeshkumar Reddy  <https://orcid.org/0000-0001-9490-2483>

Sandeep Chinta  <https://orcid.org/0000-0003-0199-1101>

Richard Matear  <https://orcid.org/0000-0002-3225-0800>

John Taylor  <https://orcid.org/0000-0001-9003-4076>

Harish Baki  <https://orcid.org/0000-0003-1956-8280>

Marcus Thatcher  <https://orcid.org/0000-0003-4139-5515>

Jatin Kala  <https://orcid.org/0000-0001-9338-2965>

Jason Sharples  <https://orcid.org/0000-0002-7816-6989>

References

- Abram N J, Henley B J, Sen Gupta A, Lippmann T J R, Clarke H, Dowdy A J, Sharples J J, Nolan R H, Zhang T and Wooster M J 2021 Connections of climate change and variability to large and extreme forest fires in southeast Australia *Commun. Earth Environ.* **2** 8
- Baki H, Chinta S, Balaji C and Srinivasan B 2022a Parameter calibration to improve the prediction of tropical cyclones over the Bay of Bengal using machine learning-based multiobjective optimization *J. Appl. Meteorol. Climatol.* **61** 819–37
- Baki H, Chinta S, Balaji C and Srinivasan B 2022b Determining the sensitive parameters of the weather research and forecasting (WRF) model for the simulation of tropical cyclones in the Bay of Bengal using global sensitivity analysis and machine learning *Geosci. Model. Dev.* **15** 2133–55
- Bjerknes V 1910 *Dynamic Meteorology and Hydrography: Part [1]-2, [and Atlas of Plates]* (Carnegie Institution of Washington)
- Bureau of Meteorology, CSIRO 2020 State of the Climate 2020

- Canadell J G, Meyer C P, Cook G D, Dowdy A, Briggs P R, Knauer J, Pepler A and Haverd V 2021 Multi-decadal increase of forest burned area in Australia is linked to climate change *Nat. Commun.* **12** 6921
- Chen F and Dudhia J 2001 Coupling an advanced land surface–hydrology model with the Penn State–NCAR MM5 modeling system part I: model implementation and sensitivity *Mon. Weather Rev.* **129** 569–85
- Chinta S and Balaji C 2020 Calibration of WRF model parameters using multiobjective adaptive surrogate model-based optimization to improve the prediction of the Indian summer monsoon *Clim. Dyn.* **55** 631–50
- Chinta S, Yaswanth Sai J and Balaji C 2021 Assessment of WRF model parameter sensitivity for high-intensity precipitation events during the Indian summer monsoon *Earth Space Sci.* **8** e2020EA001471
- Di Virgilio G, Evans J P, Blake S A P, Armstrong M, Dowdy A J, Sharples J and McRae R 2019a Climate change increases the potential for extreme wildfires *Geophys. Res. Lett.* **46** 8517–26
- Di Virgilio G, Evans J P, Di Luca A, Olson R, Argüeso D, Kala J, Andrys J, Hoffmann P, Katzfey J J and Rockel B 2019b Evaluating reanalysis-driven CORDEX regional climate models over Australia: model performance and errors *Clim. Dyn.* **53** 2985–3005
- Di Z, Duan Q, Gong W, Wang C, Gan Y, Quan J, Li J, Miao C, Ye A and Tong C 2015 Assessing WRF model parameter sensitivity: a case study with 5 day summer precipitation forecasting in the Greater Beijing Area *Geophys. Res. Lett.* **42** 579–87
- Di Z, Duan Q, Gong W, Ye A and Miao C 2017 Parametric sensitivity analysis of precipitation and temperature based on multi-uncertainty quantification methods in the weather research and forecasting model *Sci. China Earth Sci.* **60** 876–98
- Di Z, Duan Q, Wang C, Ye A, Miao C and Gong W 2018 Assessing the applicability of WRF optimal parameters under the different precipitation simulations in the Greater Beijing Area *Clim. Dyn.* **50** 1927–48
- Duan Q et al 2017 Automatic model calibration: a new way to improve numerical weather forecasting *Bull. Am. Meteorol. Soc.* **98** 959–70
- Dudhia J 1989 Numerical study of convection observed during the winter monsoon experiment using a mesoscale two-dimensional model *J. Atmos. Sci.* **46** 3077–107
- Engel C B, Lane T P, Reeder M J and Reznym M 2013 The meteorology of black saturday *Q. J. R. Meteorol. Soc.* **139** 585–99
- Evans J P, Ekström M and Ji F 2012 Evaluating the performance of a WRF physics ensemble over Southeast Australia *Clim. Dyn.* **39** 1241–58
- Evans J P, Ji F, Lee C, Smith P, Argüeso D and Fita L 2014 Design of a regional climate modelling projection ensemble experiment—NARClIM *Geosci. Model. Dev.* **7** 621–9
- Evans J P and McCabe M F 2010 Regional climate simulation over Australia's Murray-Darling basin: a multitemporal assessment *J. Geophys. Res.: Atmos.* **115** D14
- Fonseca R, Zorzano-Mier M-P, Azua-Bustos A, González-Silva C and Martín-Torres J 2019 A surface temperature and moisture intercomparison study of the weather research and forecasting model, in-situ measurements and satellite observations over the Atacama Desert *Q. J. R. Meteorol. Soc.* **145** 2202–20
- Hersbach H et al 2020 The ERA5 global reanalysis *Q. J. R. Meteorol. Soc.* **146** 1999–2049
- Hong S-Y, Dudhia J and Chen S-H 2004 A revised approach to ice microphysical processes for the bulk parameterization of clouds and precipitation *Mon. Weather Rev.* **132** 103–20
- Hong S-Y, Noh Y and Dudhia J 2006 A new vertical diffusion package with an explicit treatment of entrainment processes *Mon. Weather Rev.* **134** 2318–41
- Hong S and Lim J-O 2006 The WRF single-moment 6-Class microphysics scheme (WSM6) *Asia Pac. J. Atmos. Sci.* **42** 129–51
- Ji D, Dong W, Hong T, Dai T, Zheng Z, Yang S and Zhu X 2018 Assessing parameter importance of the weather research and forecasting model based on global sensitivity analysis methods *J. Geophys. Res.: Atmos.* **123** 4443–60
- Ji F, Nishant N, Evans J P, Di Virgilio G, Cheung K K W, Tam E, Beyer K and Riley M L 2022 Introducing NARClIM1.5: evaluation and projection of climate extremes for southeast Australia *Weather Clim. Extrem.* **38** 100526
- Jiménez P A, Dudhia J, González-Rouco J F, Navarro J, Montávez J P and García-Bustamante E 2012 A revised scheme for the WRF surface layer formulation *Mon. Weather Rev.* **140** 898–918
- Kain J S 2004 The Kain–Fritsch convective parameterization: an update *J. Appl. Meteorol.* **43** 170–81
- Kala J, Evans J P and Pitman A J 2015 Influence of antecedent soil moisture conditions on the synoptic meteorology of the Black Saturday bushfire event in southeast Australia *Q. J. R. Meteorol. Soc.* **141** 3118–29
- Kala J, Valmassoi A and Hirsch A L 2023 Assessing the maximum potential cooling benefits of irrigation in Australia during the 'Angry Summer' of 2012/2013 *Weather Clim. Extrem.* **39** 100538
- Masson-Delmotte V, Zhai P, Pirani A, Connors S L, Péan C, Berger S, Caud N, Chen Y, Goldfarb L and Gomis M I 2021 Climate change 2021: the physical science basis *Contribution of Working Group I to the Sixth Assessment Report of the Intergovernmental Panel on Climate Change* p 2
- Mlawer E J, Taubman S J, Brown P D, Iacono M J and Clough S A 1997 Radiative transfer for inhomogeneous atmospheres: RRTM, a validated correlated-k model for the longwave *J. Geophys. Res.: Atmos.* **102** 16663–82
- Montornès Torrecillas A, Codina B and Zack J W 2015 A discussion about the role of the shortwave schemes on real WRF-ARW simulations. Two case studies: cloudless and cloudy sky *Tethys: J. Mediterr. Meteorol. Climatol.* **12** 13–31
- Oke T R 2002 *Boundary Layer Climates* 2nd edn (Routledge)
- Perkins-Kirkpatrick S E and Lewis S C 2020 Increasing trends in regional heatwaves *Nat. Commun.* **11** 3357
- Quan J, Di Z, Duan Q, Gong W, Wang C, Gan Y, Ye A and Miao C 2016 An evaluation of parametric sensitivities of different meteorological variables simulated by the WRF model *Q. J. R. Meteorol. Soc.* **142** 2925–34
- Rakovec O, Hill M C, Clark M P, Weerts A H, Teuling A J and Uijlenhoet R 2014 Distributed Evaluation of Local Sensitivity Analysis (DELSA), with application to hydrologic models *Water Resour. Res.* **50** 409–26
- Reddy P J, Perkins-Kirkpatrick S E and Sharples J J 2021 Intensifying Australian heatwave trends and their sensitivity to observational data *Earths Future* **9** e2020EF001924
- Saltelli A 2002 Sensitivity analysis for importance assessment *Risk Anal.* **22** 579–90
- Saltelli A, Aleksankina K, Becker W, Fennell P, Ferretti F, Holst N, Li S and Wu Q 2019 Why so many published sensitivity analyses are false: a systematic review of sensitivity analysis practices *Environ. Modelling Softw.* **114** 29–39
- Saltelli A, Annoni P, Azzini I, Campolongo F, Ratto M and Tarantola S 2010 Variance based sensitivity analysis of model output *design and estimator for the total sensitivity index Comput. Phys. Commun.* **181** 259–70
- Sharples J J, Cary G J, Fox-Hughes P, Mooney S, Evans J P, Fletcher M-S, Fromm M, Grierson P F, McRae R and Baker P 2016 Natural hazards in Australia: extreme bushfire *Clim. Change* **139** 85–99
- Skamarock W C, Klemp J B, Dudhia J, Gill D O, Liu Z, Berner J and Huang X-yu 2021 A Description of the Advanced Research WRF Model Version 4.3 (No. NCAR/TN-556+STR) (<https://doi.org/10.5065/1dfh-6p97>)

- Sobol' I M 2001 Global sensitivity indices for nonlinear mathematical models and their Monte Carlo estimates *Math. Comput. Simul.* **55** 271–80
- Su C-H et al 2022 BARRA2: development of the next-generation Australian regional atmospheric reanalysis (available at: www.bom.gov.au/research/publications/researchreports/BRR-067.pdf)
- Temimi M et al 2020 Assessing the impact of changes in land surface conditions on WRF predictions in arid regions *J. Hydrometeorol.* **21** 2829–53
- Tewari M, Chen F, Wang W, Dudhia J, LeMone M A, Mitchell K, Ek M, Gayno G, Wegiel J and Cuenca R H, 2004 Implementation and verification of the unified NOAA land surface model in the WRF model *20th Conf. on Weather Analysis and Forecasting/16th Conf. on Numerical Weather Prediction* pp 2165–70
- Wang C et al 2020 Assessing the sensitivity of land-atmosphere coupling strength to boundary and surface layer parameters in the WRF model over Amazon *Atmos. Res.* **234** 104738
- Williams C K I and Rasmussen C E 2006 *Gaussian Processes for Machine Learning* (MIT Press)
- Williams C and Rasmussen C 1995 Gaussian processes for regression *Advances in Neural Information Processing Systems* p 8
- Yang B, Berg L K, Qian Y, Wang C, Hou Z, Liu Y, Shin H H, Hong S and Pekour M 2019 Parametric and structural sensitivities of turbine-height wind speeds in the boundary layer parameterizations in the weather research and forecasting model *J. Geophys. Res.: Atmos.* **124** 5951–69
- Yang B, Qian Y, Lin G, Leung R and Zhang Y 2012 Some issues in uncertainty quantification and parameter tuning: a case study of convective parameterization scheme in the WRF regional climate model *Atmos. Chem. Phys.* **12** 2409–27