

Substrate-Specific Evolution of Amine Dehydrogenases for Accessing Structurally Diverse Enantiopure (R)- β -Amino Alcohols

Yin, Xinjian; Gong, Wenzhong; Zeng, Yujing; Qiu, Hulin; Liu, Lan; Hollmann, Frank; Chen, Bishuang

DOI

[10.1021/acscatal.3c04995](https://doi.org/10.1021/acscatal.3c04995)

Publication date

2024

Document Version

Final published version

Published in

ACS Catalysis

Citation (APA)

Yin, X., Gong, W., Zeng, Y., Qiu, H., Liu, L., Hollmann, F., & Chen, B. (2024). Substrate-Specific Evolution of Amine Dehydrogenases for Accessing Structurally Diverse Enantiopure (R)- β -Amino Alcohols. *ACS Catalysis*, 14(2), 837-845. <https://doi.org/10.1021/acscatal.3c04995>

Important note

To cite this publication, please use the final published version (if applicable). Please check the document version above.


Copyright

Other than for strictly personal use, it is not permitted to download, forward or distribute the text or part of it, without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license such as Creative Commons.

Takedown policy

Please contact us and provide details if you believe this document breaches copyrights. We will remove access to the work immediately and investigate your claim.

Scheme 1. Kinetic Resolution and Deracemization Process for the Synthesis of (*R*)- β -Amino Alcohols



faces considerable challenges. To the best of our knowledge, no successful case has been reported yet, primarily due to two significant hurdles encountered in the utilization of amine dehydrogenase-catalyzed oxidative deamination.^{43,44} First, currently available (wild-type and engineered ones) amine dehydrogenases have been primarily developed for their reductive amination activity, resulting in relatively low catalytic efficiency for the oxidative deamination reaction,^{31,37,39,45,46} especially concerning β -amino alcohols (<10 U/g).³⁶ Additionally, the strict substrate specificity inherited from amino acid dehydrogenases also extends to the derived amine dehydrogenases, creating a constraint on the range of substrates that can be converted.^{47,48} Despite extensive research to expand the substrate spectra of amine dehydrogenases, in most cases, this remains a rather intricate issue.^{24–27,30,49,50} This is because almost all mutants are screened based on one or a limited set of substrates, and their enhanced activity often lacks generality. Consequently, when faced with novel substrates, the screening process for suitable mutants remains time-consuming and labor-intensive, even for enzymes that have been extensively studied.

In this work, we report an efficacious solution to develop engineered amine dehydrogenases exhibiting remarkable oxidative deamination activity toward structurally diverse β -amino alcohols. These engineered amine dehydrogenases, termed as β -amino alcohol dehydrogenases (β -AADHs), have demonstrated their utility in kinetic resolution or deracemization processes, yielding diverse enantiopure (*R*)- β -amino alcohols with good ee values and conversion.

RESULTS AND DISCUSSION

Access to the Oxidative Deamination Activity of β -AADHs. As starting point for our investigation, we chose a valine dehydrogenase previously identified from a hot spring metagenomic library (HsValDH3).⁵¹ Due to its considerable stability, HsValDH3 is an attractive starting point for evolution

and future application. wt-HsValDH3 does not exhibit sizable activity toward β -amino alcohols, which is why we first aimed at addressing this issue. Since a crystal structure of HsValDH3 is not available yet, we employed AlfaFold2⁵² to construct a three-dimensional model of HsValDH3. Subsequently, molecular docking with its native substrate, L-valine, was conducted to scrutinize the binding interactions (Figure 1A). According to the model, L-valine occupies a specific binding site nestled within the crevice formed between the substrate-binding domain and the coenzyme-binding domain of HsValDH3 (Figure 1A).

Utilizing insights from the resolved crystal structure and catalytic mechanism of amino acid dehydrogenases,^{53,54} we partitioned the substrate-binding pocket of HsValDH3 into three discrete regions (Figure 1B): the catalytic site, the carboxyl-binding site, and the side-chain binding site. The catalytic site encompasses residues Lys80 and Asp115, both of which are pivotal for the catalytic function of amino acid dehydrogenases (Figure 1C).^{53,54} The carboxyl-binding site is constituted by residues Asn261 and Lys68, governing the specificity for α -keto acids by forming hydrogen bonds and electrostatic interactions with the carboxyl group of L-valine (Figure 1C). Therefore, we targeted these two amino acid residues using a combinatorial saturation mutagenesis strategy, K68X/N261X (NNK degenerate codons). It is worth emphasizing here that, in contrast to previous studies,^{28,31,37,39,45,46} we aimed at oxidative deamination, for which so far no efficient high-throughput screening method was available. Fortunately, the deamination product (i.e., the β -keto alcohol) proved to be susceptible to spontaneous oxidation by tetrazolium red (TTC, Figure 1D), which enabled us to establish a semiquantitative colorimetric activity assay based on the formation of the intensely red-colored triphenylformazan (TPF) (Figure S1). This way, we identified three mutants (N261M, N261L, and N261F), exhibiting oxidative deamination activity toward 2-aminobutanol. Interestingly, N261M (M0) proved to be a better mutant for oxidative reaction than the extensively documented K68S/N261L (2M), while 2M shows superiority in the reductive direction (Figure 1E). This observation reinforces the necessity of our dedicated effort in conducting oxidative deamination-specific screening.

To gain further insight into the molecular basis of these differences, we performed a molecular docking analysis of (*S*)-2-aminobutanol and 1-hydroxy-2-butanone for M0 and 2M, respectively (Figure 1F). In the case of M0, hydrogen-bonding interactions between the α -hydroxyl group of the amino alcohol and Lys68 were observed, which positioned the starting material in a near-attack conformation to the cofactor (NAD⁺). No such interaction was observed in 2M. Rather, a new hydrogen bond network between the amino group of (*S*)-2-aminobutanol and K80 and D115 was observed, which may impede their involvement in the catalytic cycle and therefore may account for the drastically reduced catalytic activity in the oxidation of (*S*)-2-aminobutanol. Indeed, the kinetic parameters revealed that 2M exhibited a significantly lower k_{cat} value for (*S*)-2-aminobutanol compared to M0 (1.18 vs 5.23 s⁻¹) (Table 1). However, when 2M interacted with 1-hydroxy-2-butanone, the ketone group maintained a favorable catalytic distance with the catalytic residues, measuring 2.6 Å to K80 and 3.6 Å to D115. Furthermore, this binding arrangement resulted in a higher binding energy compared to M0 (−23.43 vs −21.36 kcal mol⁻¹) in line with the increased substrate affinity (lower K_{m} value) and catalytic efficiency (higher $k_{\text{cat}} K_{\text{m}}^{-1}$) for the α -hydroxy ketone (Table 1).




Figure 1. Access to the oxidative deamination activity of β -AADHs. (A) Overview of the three-dimensional model of HsValDH3 docked with L-valine. L-Valine is shown as a stick model. (B) Close-up view of the active pocket of HsValDH3. The substrate-binding pocket is shown as a surface model, with the catalytic site, carboxyl-binding site, and side-chain binding site represented in red, yellow, and cyan, respectively. (C) Residues were within the active pocket of HsValDH3. These residues constitute the catalytic site, carboxyl-binding site, and side-chain binding site, represented in red, yellow, and cyan, respectively. Hydrogen bonds that contribute to the binding of L-valine are represented as green-colored dotted lines. (D) Principle of the colorimetric high-throughput screening method for β -AADH. In the enzymatic system, β -amino alcohols undergo conversion into the corresponding α -hydroxy ketones catalyzed by β -AADH, with NAD^+ regeneration powered by the NADH oxidase (NOX). (E) Specific activities of the identified mutants toward (S)-2-aminobutanol and 1-hydroxy-2-butanone. The specific activity was determined using purified proteins. All enzymatic assays were performed in triplicate, at least, and the averaged values are reported. (F) Docking poses of (S)-2-aminobutanol and 1-hydroxy-2-butanone with N261M (M0) and K68S/N261L (2M). Hydrogen-bonding interactions are shown and highlighted as green dotted lines, and distance between substrates' active group and catalytic residues is shown and highlighted as blue dotted lines.

Substrate-Specific Evolution of M0 To Expand the Substrate Scope. The N261M mutation successfully conferred oxidative deamination activity to HsValDH3, resulting in the “first-generation” β -AADH (M0). But M0 converted only a

rather narrow range of short-chain (S)- β -amino alcohols (Figure S2). This outcome aligns with our expectations as natural amino acid dehydrogenases exhibit stringent substrate specificity, and




Figure 3. Substrate spectrum evaluation of the developed β -AADHs.

only 485 transformants, a 95% coverage was obtained.⁶¹ Compared to conventional substrate-specificity engineering strategies used for amino acid dehydrogenases or amine dehydrogenases, such as rational site-directed mutagenesis^{27,50,62,63} and iterative saturation mutagenesis (ISM),^{29,31,64} our approach exhibits several significant benefits: first, the lower screening effort enables screening for more substrates in shorter times. Second, it allows the capture of potential synergistic effects arising from the combination of mutations at the five selected sites, an outcome that is nearly impossible to achieve by using traditional engineering strategies.

As shown in Figure 2C, for all 7 model substrates screened, several mutants with significantly increased catalytic activity were observed. The best-performing mutants for *L*-valinol, (*S*)-2-amino-4-methylpentane-1-ol, and (*S*)-2-aminohexane-1-ol exhibited a specific activity improvement of 1.5- to 27.6-fold compared to that of M0. Furthermore, for the four substrates initially showing no activity with M0, the generated mutants acquired substantial catalytic activity (0.15–1.22 U/mg).

Particularly mutants M1–M4 (M1: E114G/N261M; M2: A113G/E114A/N261M; M3: A113G/E114A/P146A/N261M; M4: A113G/E114A/P146A/N261M/I294G) excelled in their catalytic performance.

Next, we examined the substrate-binding pocket structure of the best mutants using POVME 3.0.⁶⁵ A compelling correlation between the active pocket size and substrate specificity (Figure 2D) was observed. As the side chains of the substrates used for screening increased, the pocket size of the best mutants obtained also gradually increased, indicating a notable positive correlation. It serves to validate the effectiveness and efficacy of our phylogenetic analysis-guided double-code saturation mutagenesis strategy in obtaining individual optimal mutants for each substrate.

Applications of Developed β -AADHs. To assess the applicability of the developed β -AADHs, we conducted a systematic evaluation of their substrate spectra. As illustrated in

Figure 3, the “second-generation” β -AADHs (M1–M4) exhibited a remarkably expanded substrate scope compared to M0. Catalytic activity was observed for essentially all substrates tested.

Next, we explored the synthetic usefulness of newly generated β -AADHs. Six representative, racemic β -amino alcohol substrates (S1, S2, S3, S4, S9, and S12) were tested first in a kinetic resolution-type reaction using the previously identified optimal β -AADH mutant (Figure 3). For cofactor regeneration, the recombinant NADH oxidase (NOX) from *Streptococcus equi* was used.⁶⁶ As shown in Figure 4A, the majority of these model compounds were smoothly oxidized in high enantioselectivity, resulting in 50% conversion (i.e., full theoretical) and essentially optically pure (*R*)- β -amino alcohols. In two cases, even after prolonged reaction times, the conversions did not exceed 49%, and consequently, the optical purities of the remaining amino alcohols were disappointingly low (92–96% ee). The origin of this slower reaction rate is not fully understood and will be investigated in future studies. Possibly, low affinity toward the starting material (high K_m values), product inhibition, and/or enzyme inactivation may account for this.

Inspired by previous work regarding deracemization of racemic amines without α -functionalization,³³ we also evaluated the deracemization of the racemic β -amino alcohols by extending the reaction shown in Figure 4A with an enantioselective transamination of the undesired keto alcohol byproduct (Figure 4B). For this, we chose the (*R*)-selective ω -transaminase from *Bacillus megaterium* (BmTA) (Table S1).⁶⁷ The stereoselectivity of available (*R*)-selective ω -transaminases is generally low.^{19,68,69} This implies that the amino alcohols produced by these transaminases will not be enantiomerically pure. However, the proposed three-enzyme cascade deracemization system effectively addresses this issue as undesired (*S*)- β -amino alcohols are transferred back to α -hydroxy ketones by the engineered β -AADHs. Consequently, only the (*R*)-product is preserved in the system, theoretically giving excellent optical




Figure 4. Kinetic resolution and deracemization process for the synthesis of (*R*)- β -amino alcohols. (A) Kinetic resolution of racemic β -amino alcohols. Reaction conditions: β -AADHs (2 mg/mL purified protein), NOX (2 mg/mL purified protein), NAD^+ (0.5 mM), racemic β -amino alcohols (10 mM), pH 9.0 Tris-HCl (0.1 M), 2 mL of total volume. The reaction was performed at 37 °C and 200 rpm for 24 or 48 h. Conversion for kinetic resolution process was defined as the percentage ratio of consumed (*S*)- β -amino alcohols to the initial racemic β -amino alcohol substrate. (B) Deracemization of racemic β -amino alcohols (10 mM). Reaction conditions: β -AADHs (2 mg/mL purified protein), NOX (2 mg/mL purified protein), *BmTA* (2 mg/mL for S1, S2, S3, S4; 10 mg/mL for S9 and S12, purified protein), NAD^+ (0.5 mM), PLP (0.5 mM), racemic β -amino alcohols (10 mM), (*S*)-phenethylamine (50 mM), pH 9.0 Tris-HCl (0.1 M), 2 mL of total volume. The reaction was performed at 37 °C, 200 rpm for 24 or 48 h. Conversion for the deracemization process was defined as the percentage ratio of the (*R*)- β -amino alcohol product to the initial racemic β -amino alcohol substrate. (C–E) Deracemization of racemic S2, S4, and S9 (25 or 50 mM). Reaction conditions: NAD^+ (0.5 mM), PLP (0.5 mM), racemic β -amino alcohols (25 or 50 mM), and (*S*)-phenethylamine (100 mM), pH 9.0 Tris-HCl (0.1 M), 2 mL of total volume. The enzymes [1 mg/mL β -AADHs (M1 for S2, M2 for S4, and M3 for S9), 2 mg/mL NOX, and 2 mg/mL *BmTA*] were added into the reaction system at specific time points: 0, 6, 12, 24, and 36 h. The reaction was performed at 37 °C, 200 rpm.

purity for various β -amino alcohol products. Gratifyingly, all six racemic β -amino alcohols were efficiently deracemized, resulting in the production of (*R*)- β -amino alcohols with ee values >99% and conversions ranging from 90 to 99%. Even for substrates S3 and S4, which exhibited imperfect ee values in the kinetic resolution process, the deracemization process yielded satisfactory results.

Motivated by these results, we proceeded to explore the deracemization process with higher substrate loadings (25–50 mM) (Figure 4C–E). To surmount enzyme deactivation issues observed in prior batch reactions (Figure S5), particularly concerning NOX and *BmTA*, we added these enzymes at intervals (0, 6, 12, 24, and 36 h). Applying this strategy, complete deracemization of 25 mM racemic S2, S4, and S9 substrates was achieved within 10, 30, and 36 h, respectively, with optical purities surpassing 99% ee. Upon further increasing the substrate concentration to 50 mM, S2 and S4 achieved complete deracemization within 30 and 48 h (ee > 99%), respectively, while the final ee value for S9 stood at 76%.

Preparative-scale deracemization reactions were performed on a 50 mL scale with 50 mM racemic S2 and S4. These

reactions mirror the process observed in the 2 mL-scale reactions (Figure S6). The deracemization of racemic S2 and S4 reached completion (>99% conversion) within 34 and 48 h (ee > 99%), respectively, yielding a final concentration of (*R*)-R2 and (*R*)-R4.

Compared with extensively explored and optimized amine dehydrogenase-catalyzed reductive amination processes for (*S*)- β -amino alcohol production,^{37–39} the productivity observed here (2.1 mM h⁻¹) somewhat falls back (up to 100 mM h⁻¹) albeit also using approximately 20–100 times less of the biocatalyst. Hence, we are convinced that further reaction engineering will yield reaction schemes at least comparable to those of the above-mentioned (*S*)-amino alcohol syntheses.

CONCLUSIONS

In this contribution, we have expanded the biocatalytic toolbox for chiral β -amino alcohol synthesis with a newly designed amine dehydrogenase from a wild-type valine dehydrogenase. A mutation N261M in the carboxyl-binding site was identified to endow HsValDH3 with oxidative deamination activity, creating the “first-generation” β -AADH (M0). Guided by phylogenetic

analysis, double-code saturation mutagenesis in MO's substrate-binding pocket led to the "second-generation" β -AADHs (M1–M4) with broadened substrate spectra.

Also, the newly developed β -AADHs themselves exhibit a significant potential for the preparative kinetic resolution and deracemization of readily available racemic β -amino alcohols.

Further work in our laboratory will focus on the further optimization of the methodology, generating even more active and selective β -AADHs and embedding the deracemization in more complex cascades to convert simple starting materials into diverse value-added products.

■ ASSOCIATED CONTENT

SI Supporting Information

The Supporting Information is available free of charge at <https://pubs.acs.org/doi/10.1021/acscatal.3c04995>.

Detailed description of experimental procedures; analytic conditions; primer sequences; enzyme sequences; GC conditions; HPLC conditions; colorimetric high-throughput screening results; specific activities of amino acid dehydrogenases; Michaelis–Menten diagrams; pre-column derivatization HPLC analysis; GC analysis; sequence information; and supporting HPLC, NMR, and MS spectra (PDF)

■ AUTHOR INFORMATION

Corresponding Authors

Frank Hollmann – Department of Biotechnology, Delft University of Technology, Delft 2629HZ, The Netherlands; orcid.org/0000-0003-4821-756X; Email: f.hollmann@tudelft.nl

Bishuang Chen – School of Marine Sciences, Sun Yat-Sen University, Zhuhai 519080, China; Southern Marine Science and Engineering Guangdong Laboratory (Zhuhai), Zhuhai 519080, China; orcid.org/0000-0002-7221-1208; Email: chenbsh23@mail.sysu.edu.cn

Authors

Xinjian Yin – School of Marine Sciences, Sun Yat-Sen University, Zhuhai 519080, China; Southern Marine Science and Engineering Guangdong Laboratory (Zhuhai), Zhuhai 519080, China

Wenzhong Gong – School of Marine Sciences, Sun Yat-Sen University, Zhuhai 519080, China

Yujing Zeng – School of Marine Sciences, Sun Yat-Sen University, Zhuhai 519080, China

Hulin Qiu – School of Marine Sciences, Sun Yat-Sen University, Zhuhai 519080, China

Lan Liu – School of Marine Sciences, Sun Yat-Sen University, Zhuhai 519080, China; Southern Marine Science and Engineering Guangdong Laboratory (Zhuhai), Zhuhai 519080, China

Complete contact information is available at: <https://pubs.acs.org/doi/10.1021/acscatal.3c04995>

Author Contributions

The manuscript was written through contributions of all authors. All authors have given approval to the final version of the manuscript.

Notes

The authors declare no competing financial interest.

■ ACKNOWLEDGMENTS

This work was financially supported by the National Natural Science Foundation of China (Nos. 22108317 and 42376097); Guangdong Basic and Applied Basic Research Foundation (2019A1515111051, 2021A1515010829, and 2023A1515030226); and China Postdoctoral Science Foundation (2020M680134).

■ REFERENCES

- (1) Zhou, Z. J.; Tan, Y. Q.; Shen, X.; Ivlev, S.; Meggers, E. Catalytic enantioselective synthesis of β -amino alcohols by nitrene insertion. *Sci. China Chem.* **2021**, *64*, 452–458.
- (2) Nakafuku, K. M.; Zhang, Z.; Wappes, E. A.; Stateman, L. M.; Chen, A. D.; Nagib, D. A. Enantioselective radical C–H amination for the synthesis of β -amino alcohols. *Nat. Chem.* **2020**, *12*, 697–704.
- (3) Reddy, U. V. S.; Chennapuram, M.; Seki, C.; Kwon, E.; Okuyama, Y.; Nakano, H. Catalytic Efficiency of Primary β -Amino Alcohols and Their Derivatives in Organocatalysis. *Eur. J. Org. Chem.* **2016**, *2016*, 4124–4143.
- (4) Vicario, J. L.; Badía, D.; Carrillo, L.; Reyes, E.; Etxebarria, J. α -Amino acids, β -amino alcohols and related compounds as chiral auxiliaries, ligands and catalysts in the asymmetric aldol reaction. *Curr. Org. Chem.* **2005**, *9*, 219–235.
- (5) Bergmeier, S. C. The synthesis of vicinal amino alcohols. *Tetrahedron* **2000**, *56*, 2561–2576.
- (6) Ager, D. J.; Prakash, I.; Schaad, D. R. 1,2-Amino alcohols and their heterocyclic derivatives as chiral auxiliaries in asymmetric synthesis. *Chem. Rev.* **1996**, *96*, 835–875.
- (7) Demange, L.; Lozach, O.; Ferandin, Y.; Hoang, N. T.; Meijer, L.; Galons, H. Synthesis and evaluation of new potent inhibitors of CK1 and CDK5, two kinases involved in Alzheimer's disease. *Med. Chem. Res.* **2013**, *22*, 3247–3258.
- (8) Oumata, N.; Bettayeb, K.; Ferandin, Y.; Demange, L.; Lopez-Giral, A.; Goddard, M.-L.; Myriantopoulos, V.; Mikros, E.; Flajolet, M.; Greengard, P.; Meijer, L.; Galons, H. Roscovitine-derived, dual-specificity inhibitors of cyclin-dependent kinases and casein kinases 1. *J. Med. Chem.* **2008**, *51* (17), 5229–5242.
- (9) Yuan, S.; Yu, B.; Liu, H.-M. New drug approvals for 2019: Synthesis and clinical applications. *Eur. J. Med. Chem.* **2020**, *205*, No. 112667.
- (10) Karlström, S.; Nordvall, G.; Sohn, D.; Hettman, A.; Turek, D.; Åhlin, K.; Kers, A.; Claesson, M.; Slivo, C.; Lo-Alfredsson, Y.; Petersson, C.; Bessidskaia, G.; Svensson, P. H.; Rein, T.; Jerning, E.; Malmberg, A.; Ahlgen, C.; Ray, C.; Vares, R.; Ivanov, V.; Johansson, R. Substituted 7-amino-5-thio-thiazolo [4, 5-d] pyrimidines as potent and selective antagonists of the fractalkine receptor (CX3CR1). *J. Med. Chem.* **2013**, *56* (8), 3177–3190.
- (11) Hu, H.; Wang, Z. B. Cr-Catalyzed Asymmetric Cross Aza-Pinacol Couplings for β -Amino Alcohol Synthesis. *J. Am. Chem. Soc.* **2023**, *145*, 20775–20781.
- (12) Azizi, N.; Saidi, M. R. Highly Chemoselective Addition of Amines to Epoxides in Water. *Org. Lett.* **2005**, *7*, 3649–3651.
- (13) Azoulay, S.; Manabe, K.; Kobayashi, S. Catalytic Asymmetric Ring Opening of meso-Epoxides with Aromatic Amines in Water. *Org. Lett.* **2005**, *7*, 4593–4595.
- (14) Liu, B. X.; Xie, P. F.; Zhao, J.; Wang, J. J.; Wang, M. M.; Jiang, Y. Q.; Chang, J. B.; Li, X. W. Rhodium-Catalyzed Enantioselective Synthesis of β -Amino Alcohols via Desymmetrization of gem-Dimethyl Groups. *Angew. Chem., Int. Ed.* **2021**, *60*, 8396–8400.
- (15) Shivani; Pujala, B.; Chakraborti, A. K. Zinc(II) Perchlorate Hexahydrate Catalyzed Opening of Epoxide Ring by Amines: Applications to Synthesis of (R)/(R)-Propranolols and (R)/(R)/(S)-Naftopidils. *J. Org. Chem.* **2007**, *72*, 3713–3722.
- (16) Gupta, P.; Mahajan, N. Biocatalytic approaches towards the stereoselective synthesis of vicinal amino alcohols. *New J. Chem.* **2018**, *42*, 12296–12327.
- (17) Pavlidis, I. V.; Weiß, M. S.; Genz, M.; Spurr, P.; Hanlon, S. P.; Wirz, B.; Iding, H.; Bornscheuer, U. T. Identification of (S)-selective

- transaminases for the asymmetric synthesis of bulky chiral amines. *Nat. Chem.* **2016**, *8*, 1076–1082.
- (18) Sekar, B. S.; Mao, J.; Lukito, B. R.; Wang, Z.; Li, Z. Bioproduction of Enantiopure (R)- and (S)-2-Phenylglycinols from Styrenes and Renewable Feedstocks. *Adv. Synth. Catal.* **2021**, *363*, 1892–1903.
- (19) Zhang, J. D.; Yang, X. X.; Dong, R.; Gao, L. L.; Li, J.; Li, X.; Huang, S. P.; Zhang, C. F.; Chang, H. H. Cascade Biocatalysis for Regio- and Stereoselective Aminohydroxylation of Styrenyl Olefins to Enantiopure Arylglycinols. *ACS Sus. Chem. Eng.* **2020**, *8*, 18277–18285.
- (20) Zhang, J. D.; Yang, X. X.; Jia, Q.; Zhao, J. W.; Gao, L. L.; Gao, W. C.; Chang, H. H.; Wei, W. L.; Xu, J. H. Asymmetric ring opening of racemic epoxides for enantioselective synthesis of (S)- β -amino alcohols by a cofactor self-sufficient cascade biocatalysis system. *Catal. Sci. Technol.* **2019**, *9*, 70–74.
- (21) Corrado, M. L.; Knaus, T.; Mutti, F. G. Regio- and stereoselective multi-enzymatic aminohydroxylation of β -methylstyrene using dioxygen, ammonia and formate. *Green Chem.* **2019**, *21*, 6246–6251.
- (22) Corrado, M. L.; Knaus, T.; Schwaneberg, U.; Mutti, F. G. High-Yield Synthesis of Enantiopure 1,2-Amino Alcohols from 1-Phenylalanine via Linear and Divergent Enzymatic Cascades. *Org. Process Res. Dev.* **2022**, *26*, 2085–2095.
- (23) González-Granda, S.; Steinkellner, G.; Gruber, K.; Lavandera, L.; Gotor-Fernández, V. Gold and Biocatalysis for the Stereodivergent Synthesis of Nor(pseudo)ephedrine Derivatives: Cascade Design Toward Amino Alcohols, Diols, and Diamines. *Adv. Synth. Catal.* **2023**, *365*, 1036–1047.
- (24) Pushpanath, A.; Siirola, E.; Bornadel, A.; Woodlock, D.; Schell, U. Understanding and Overcoming the Limitations of *Bacillus badius* and *Caldalkalibacillus thermarum* Amine Dehydrogenases for Biocatalytic Reductive Amination. *ACS Catal.* **2017**, *7*, 3204–3209.
- (25) Chen, F. F.; Zheng, G. W.; Liu, L.; Li, H.; Chen, Q.; Li, F. L.; Li, C. X.; Xu, J. H. Reshaping the Active Pocket of Amine Dehydrogenases for Asymmetric Synthesis of Bulky Aliphatic Amines. *ACS Catal.* **2018**, *8*, 2622–2628.
- (26) Franklin, R. D.; Mount, C. J.; Bommarius, B. R.; Bommarius, A. S. Separate Sets of Mutations Enhance Activity and Substrate Scope of Amine Dehydrogenase. *ChemCatChem.* **2020**, *12*, 2436–2439.
- (27) Kong, W. X.; Liu, Y. T.; Huang, C.; Zhou, L. Y.; Gao, J.; Turner, N. J.; Jiang, Y. J. Direct Asymmetric Reductive Amination of Alkyl (Hetero)Aryl Ketones by an Engineered Amine Dehydrogenase. *Angew. Chem., Int. Ed.* **2022**, *61*, No. e202202264.
- (28) Liu, L.; Wang, D. H.; Chen, F. F.; Zhang, Z. J.; Chen, Q.; Xu, J. H.; Wang, Z. L.; Zheng, G. W. Development of an engineered thermostable amine dehydrogenase for the synthesis of structurally diverse chiral amines. *Catal. Sci. Technol.* **2020**, *10*, 2353–2358.
- (29) Ming, H.; Yuan, B.; Qu, G.; Sun, Z. T. Engineering the activity of amine dehydrogenase in the asymmetric reductive amination of hydroxyl ketones. *Catal. Sci. Technol.* **2022**, *12*, 5952–5960.
- (30) Mu, X. Q.; Wu, T.; Mao, Y.; Zhao, Y. L.; Xu, Y.; Nie, Y. Iterative Alanine Scanning Mutagenesis Confers Aromatic Ketone Specificity and Activity of L-Amine Dehydrogenases. *ChemCatChem.* **2021**, *13*, 5243–5253.
- (31) Qian, Y. Y.; Chu, L. L.; Zhang, X. Y.; Lu, Z. W.; Bai, Y. P. Stereoselective Synthesis of Structurally Diverse (S)-Lactams via an Engineered Amine Dehydrogenase. *Adv. Synth. Catal.* **2022**, *364*, 4289–4299.
- (32) Ye, L. J.; Toh, H. H.; Yang, Y.; Adams, J. P.; Snajdrova, R.; Li, Z. Engineering of amine dehydrogenase for asymmetric reductive amination of ketone by evolving *Rhodococcus phenylalanine* dehydrogenase. *ACS Catal.* **2015**, *5*, 1119–1122.
- (33) Yoon, S.; Patil, M. D.; Sarak, S.; Jeon, H.; Kim, G. H.; Khobragade, T. P.; Sung, S.; Yun, H. Deracemization of Racemic Amines to Enantiopure (R)- and (S)-amines by Biocatalytic Cascade Employing ω -Transaminase and Amine Dehydrogenase. *ChemCatChem.* **2019**, *11*, 1898–1902.
- (34) Tseliou, V.; Knaus, T.; Vilím, J.; Masman, M. F.; Mutti, F. G. Kinetic Resolution of Racemic Primary Amines Using *Geobacillus stearothermophilus* Amine Dehydrogenase Variant. *ChemCatChem.* **2020**, *12*, 2184–2188.
- (35) Patil, M. D.; Yoon, S.; Jeon, H.; Khobragade, T. P.; Sarak, S.; Pagar, A. D.; Won, Y.; Yun, H. Kinetic resolution of racemic amines to enantiopure (S)-amines by a biocatalytic cascade employing amine dehydrogenase and alanine dehydrogenase. *Catalysts* **2019**, *9*, 600.
- (36) Jeon, H.; Yoon, S.; Ahsan, M. M.; Sung, S.; Kim, G. H.; Sundaramoorthy, U.; Rhee, S. K.; Yun, H. The kinetic resolution of racemic amines using a whole-cell biocatalyst co-expressing amine dehydrogenase and NADH oxidase. *Catalysts* **2017**, *7*, 251.
- (37) Chen, F. F.; Cosgrove, S. C.; Birmingham, W. R.; Mangas-Sanchez, J.; Citoler, J.; Thompson, M. P.; Zheng, G. W.; Xu, J. H.; Turner, N. J. Enantioselective Synthesis of Chiral Vicinal Amino Alcohols Using Amine Dehydrogenases. *ACS Catal.* **2019**, *9*, 11813–11818.
- (38) Tong, F. F.; Qin, Z. M.; Wang, H. Y.; Jiang, Y. Y.; Li, J. K.; Ming, H.; Qu, G.; Xiao, Y. Z.; Sun, Z. T. Biosynthesis of Chiral Amino Alcohols via an Engineered Amine Dehydrogenase in *E. coli*. *Front. Bioeng. Biotechnol.* **2022**, *9*, No. 778584.
- (39) Wang, H.; Qu, G.; Li, J. K.; Ma, J. A.; Guo, J.; Miao, Y.; Sun, Z. Data mining of amine dehydrogenases for the synthesis of enantiopure amino alcohols. *Catal. Sci. Technol.* **2020**, *10*, 5945–5952.
- (40) Mayol, O.; Bastard, K.; Beloti, L.; Frese, A.; Turkenburg, J. P.; Petit, J.-L.; Mariage, A.; Debard, A.; Pellouin, V.; Perret, A.; de Berardinis, V.; Zaparucha, A.; Grogan, G.; Vergne-Vaxelaire, C. A family of native amine dehydrogenases for the asymmetric reductive amination of ketones. *Nat. Catal.* **2019**, *2* (4), 324–333.
- (41) Abiko, A.; Masamune, S. An improved, convenient procedure for reduction of amino acids to aminoalcohols: Use of $\text{NaBH}_4\text{-H}_2\text{SO}_4$. *Tetrahedron Lett.* **1992**, *33*, 5517–5518.
- (42) Zhang, J. D.; Zhao, J. W.; Gao, L. L.; Zhao, J.; Chang, H. H.; Wei, W. L. One-Pot Three-Step Consecutive Transformation of L- α -Amino Acids to (R)- and (S)-Vicinal 1,2-Diols via Combined Chemical and Biocatalytic Process. *ChemCatChem.* **2019**, *11*, 5032–5037.
- (43) Knaus, T.; Böhmer, W.; Mutti, F. G. Amine dehydrogenases: Efficient biocatalysts for the reductive amination of carbonyl compounds. *Green Chem.* **2017**, *19*, 453–463.
- (44) Liu, J. Q.; Kong, W. X.; Bai, J.; Li, Y. X.; Dong, L. L.; Zhou, L. Y.; Liu, Y. T.; Gao, J.; Bradshaw Allen, R. T.; Turner, N. J.; Jiang, Y. J. Amine dehydrogenases: Current status and potential value for chiral amine synthesis. *Chem. Catal.* **2022**, *2*, 1288–1314.
- (45) Abrahamson, M. J.; Vázquez-Figueroa, E.; Woodall, N. B.; Moore, J. C.; Bommarius, A. S. Development of an amine dehydrogenase for synthesis of chiral amines. *Angew. Chem., Int. Ed.* **2012**, *51*, 3969–3972.
- (46) Abrahamson, M. J.; Wong, J. W.; Bommarius, A. S. The evolution of an amine dehydrogenase biocatalyst for the asymmetric production of chiral amines. *Adv. Synth. Catal.* **2013**, *355*, 1780–1786.
- (47) Ohshima, T.; Soda, K. Biochemistry and biotechnology of amino acid dehydrogenases. *Adv. Biochem. Eng. Biotechnol.* **1990**, *42*, 187–209.
- (48) Zhou, F.; Xu, Y.; Nie, Y.; Mu, X. Q. Substrate-Specific Engineering of Amine Dehydrogenase Superfamily for Synthesis of a Variety of Chiral Amines and Amino Acids. *Catalysts* **2022**, *12*, 380.
- (49) Bommarius, B. R.; Schürmann, M.; Bommarius, A. S. A novel chimeric amine dehydrogenase shows altered substrate specificity compared to its parent enzymes. *Chem. Commun.* **2014**, *50*, 14953–14955.
- (50) Wang, D. H.; Chen, Q.; Yin, S. N.; Ding, X. W.; Zheng, Y. C.; Zhang, Z.; Zhang, Y. H.; Chen, F. F.; Xu, J. H.; Zheng, G. W. Asymmetric Reductive Amination of Structurally Diverse Ketones with Ammonia Using a Spectrum-Extended Amine Dehydrogenase. *ACS Catal.* **2021**, *11*, 14274–14283.
- (51) Yin, X. J.; Gong, W. Z.; Zhan, Z. G.; Wei, W.; Li, M. M.; Jiao, J. Y.; Chen, B. S.; Liu, L.; Li, W. J.; Gao, Z. A. Mining and engineering of valine dehydrogenases from a hot spring sediment metagenome for the synthesis of chiral non-natural L-amino acids. *Mol. Catal.* **2022**, *533*, No. 112767.
- (52) Jumper, P.; Evans, R.; Pritzel, A.; Green, T.; Figurnov, M.; Ronneberger, O.; Tunyasuvunakool, K.; Bates, R.; Židek, A.;

Potapenko, A.; Bridgland, A.; Meyer, C.; Kohl, S. A. A.; Ballard, A. J.; Cowie, A.; Romera-Paredes, B.; Nikolov, S.; Jain, R.; Adler, J.; Back, T.; Petersen, S.; Reiman, D.; Clancy, E.; Zielinski, M.; Steinegger, M.; Pacholska, M.; Berghammer, T.; Bodenstein, S.; Silver, D.; Vinyals, O.; Senior, A. W.; Kavukcuoglu, K.; Kohli, P.; Hassabis, D. Highly accurate protein structure prediction with AlphaFold. *Nature* **2021**, *596*, 583–589.

(53) Turnbull, A. P.; Baker, P. J.; Rice, D. W. Analysis of the quaternary structure, substrate specificity, and catalytic mechanism of valine dehydrogenase. *J. Biol. Chem.* **1997**, *272*, 25105–25111.

(54) Brunhuber, N. M. W.; Thoden, J. B.; Blanchard, J. S.; Vanhooke, J. L. *Rhodococcus* L-phenylalanine dehydrogenase: Kinetics, mechanism, and structural basis for catalytic specificity. *Biochem.* **2000**, *39*, 9174–9187.

(55) Wu, T.; Mu, X. Q.; Xue, Y. Y.; Xu, Y.; Nie, Y. Structure-guided steric hindrance engineering of *Bacillus badius* phenylalanine dehydrogenase for efficient L-homophenylalanine synthesis. *Biotechnol. Biofuels* **2021**, *14*, 207.

(56) Meng, X. Q.; Yang, L.; Liu, Y.; Wang, H. L.; Shen, Y. L.; Wei, D. Z. Identification and Rational Engineering of a High Substrate-Tolerant Leucine Dehydrogenase Effective for the Synthesis of L-tert-Leucine. *ChemCatChem* **2021**, *13*, 3340–3349.

(57) Li, J.; Pan, J.; Zhang, J.; Xu, J. H. Stereoselective synthesis of L-tert-leucine by a newly cloned leucine dehydrogenase from *Exiguobacterium sibiricum*. *J. Mol. Catal. B. Enzym.* **2014**, *105*, 11–17.

(58) Asano, Y.; Yamada, A.; Kato, Y.; Yamaguchi, K.; Hibino, Y.; Hirai, K.; Kondo, K. Enantioselective Synthesis of (S)-Amino Acids by Phenylalanine Dehydrogenase from *Bacillus sphaericus*: Use of Natural and Recombinant Enzymes. *J. Org. Chem.* **1990**, *55*, 5567–5571.

(59) Schütte, H.; Hummel, W.; Tsai, H.; Kula, M. R. L-leucine dehydrogenase from *Bacillus cereus* - Production, large-scale purification and protein characterization. *Appl. Microbiol. Biotechnol.* **1985**, *22*, 306–317.

(60) Ohshima, T.; Misono, H.; Soda, K. Properties of crystalline leucine dehydrogenase from *Bacillus sphaericus*. *J. Biol. Chem.* **1978**, *253*, 5719–5725.

(61) Reetz, M. T.; Kahakeaw, D.; Lohmer, R. Addressing the Numbers Problem in Directed Evolution. *ChemBioChem* **2008**, *9*, 1797–1804.

(62) Wang, Z. Y.; Zhou, H. S.; Yu, H. R.; Pu, Z. J.; Xu, J. L.; Zhang, H. Y.; Wu, J. P.; Yang, L. R. Computational Redesign of the Substrate Binding Pocket of Glutamate Dehydrogenase for Efficient Synthesis of Noncanonical L-Amino Acids. *ACS Catal.* **2022**, *12*, 13619–13629.

(63) Wu, T.; Wang, Y. M.; Zhang, N. X.; Yin, D. J.; Xu, Y.; Nie, Y.; Mu, X. Q. Reshaping Substrate-Binding Pocket of Leucine Dehydrogenase for Bidirectionally Accessing Structurally Diverse Substrates. *ACS Catal.* **2023**, *13*, 158–168.

(64) Lv, T.; Feng, J. H.; Chen, X.; Luo, Y. Y.; Wu, Q. Q.; Zhu, D. M.; Ma, Y. H. Desymmetric Reductive Amination of 1,3-Cyclopentadiones to Single Stereoisomer of β -Amino Ketones with an All-Carbon Quaternary Stereocenter by Engineered Amine Dehydrogenases. *ACS Catal.* **2023**, *13*, 5053–5061.

(65) Wagner, J. R.; Sørensen, J.; Hensley, N.; Wong, C.; Zhu, C.; Perison, T.; Amaro, R. E. POVME 3.0: Software for Mapping Binding Pocket Flexibility. *J. Chem. Theory Comput.* **2017**, *13*, 4584–4592.

(66) Rehn, G.; Pedersen, A. T.; Woodley, J. M. Application of NAD(P)H oxidase for cofactor regeneration in dehydrogenase catalyzed oxidations. *J. Mol. Catal. B. Enzym.* **2016**, *134*, 331–339.

(67) Hanson, R. L.; Davis, B. L.; Chen, Y. J.; Goldberg, S. L.; Parker, W. L.; Tully, T. P.; Montana, M. A.; Patel, R. N. Preparation of (R)-amines from racemic amines with an (S)-amine transaminase from *Bacillus megaterium*. *Adv. Synth. Catal.* **2008**, *350*, 1367–1375.

(68) Koszelewski, D.; Göritz, M.; Clay, D.; Seisser, B.; Kroutil, W. Synthesis of optically active amines employing recombinant ω -transaminases in *E. coli* cells. *ChemCatChem* **2010**, *2*, 73–7.

(69) Wu, H. L.; Zhang, J. D.; Zhang, C. F.; Fan, X. J.; Chang, H. H.; Wei, W. L. Characterization of Four New Distinct ω -Transaminases from *Pseudomonas putida* NBRC 14164 for Kinetic Resolution of Racemic Amines and Amino Alcohols. *Appl. Biochem. Biotechnol.* **2017**, *181*, 972–985.