

Deep plumbing model of the Cenozoic Manzaz / Atakor intraplate volcanic system, Central Hoggar, Northwest Africa, based on electrical resistivity models

Boukhalfa , Zakaria ; Comeau, Matthew J.; Benhallou , Amel ; Bouzid , Abderrezak ; Bendaoud , Abderrahmane

DOI

[10.5194/egusphere-egu24-473](https://doi.org/10.5194/egusphere-egu24-473)

Publication date

2024

Document Version

Final published version

Citation (APA)

Boukhalfa , Z., Comeau, M. J., Benhallou , A., Bouzid , A., & Bendaoud , A. (2024). *Deep plumbing model of the Cenozoic Manzaz / Atakor intraplate volcanic system, Central Hoggar, Northwest Africa, based on electrical resistivity models*. Abstract from EGU General Assembly 2024, Vienna, Austria.
<https://doi.org/10.5194/egusphere-egu24-473>

Important note

To cite this publication, please use the final published version (if applicable).
Please check the document version above.

Copyright

Other than for strictly personal use, it is not permitted to download, forward or distribute the text or part of it, without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license such as Creative Commons.

Takedown policy

Please contact us and provide details if you believe this document breaches copyrights.
We will remove access to the work immediately and investigate your claim.

EGU24-473, updated on 28 Mar 2024

<https://doi.org/10.5194/egusphere-egu24-473>

EGU General Assembly 2024

© Author(s) 2024. This work is distributed under the Creative Commons Attribution 4.0 License.



Deep plumbing model of the Cenozoic Manzaz / Atakor intraplate volcanic system, Central Hoggar, Northwest Africa, based on electrical resistivity models

Zakaria Boukhalfa^{1,3}, Matthew J. Comeau², Amel Benhallou¹, Abderrezak Bouzid¹, and Abderrahmane Bendaoud³

¹Centre de Recherche en Astronomie, Astrophysique et Géophysique (CRAAG), Subsurface Geophysics, Algiers, Algeria (boukhalfa.zakaria@yahoo.com)

²Delft University of Technology, Department of Geoscience and Engineering, Delft, The Netherlands.

³Faculté des sciences de la terre, de Géographie et de l'Aménagement du Territoire (FSTGAT), Université des Sciences et Technologie Houari Boumediene (USTHB), Algiers, Algeria.

Continental intraplate volcanic systems, with their location far from plate tectonic boundaries, are not well understood: the crustal and lithospheric mantle structure of these systems remain enigmatic and there is no consensus on the mechanisms that cause melt generation and ascent. The Cenozoic saw the development of numerous volcanic provinces on the African plate. This includes the Hoggar volcanic province, located in Northwest Africa, part of the Tuareg shield. It is composed of several massifs with contrasting ages and eruptive styles. The magmatic activity began at around 34 Ma and continued throughout the Neogene-Quaternary. Phonolite and trachyte domes as well as scoria cones and necks are found in the Manzaz and Atakor volcanic districts. In order to image the crustal and lithospheric mantle structure of this region, and to understand the origins and potential mechanisms of the continental intraplate volcanic activity in the Central Hoggar and specifically the Atakor/Manzaz area, we acquired magnetotelluric (MT) measurements from 40 locations and generated a 3-D electrical resistivity model. The model covers an area of about 100 km by 200 km. Images of the subsurface architecture, in terms of electrical resistivity, from the near-surface to the lithospheric mantle, allow us to image the deep plumbing system of the volcanic system. Low resistivity features (i.e., conductors) in the crust that are narrow, linear structures trending approximately north-south, are revealed along the two boundaries of the Azrou N'Fad terrane, in the Manzaz area. They likely reflect the Pan-African mega-shear zones, which were reactivated throughout the tectonic evolution of the region. The model reveals that these faults are lithospheric-scale. In addition, the low-resistivity features likely represent the signatures of past fluid flow. The location of the recent Cenozoic volcanic activity was likely influenced by the pre-existing structure. A deep feature of moderate conductivity is located in the upper lithospheric mantle directly beneath the Manzaz and Atakor Volcanic Districts. It may represent the origin of the overlying anomalies and may suggest metasomatism of the sub-continental lithospheric mantle.

Keywords: intraplate, Hoggar, alkaline volcanism, magnetotelluric, electrical resistivity.